

Sabouraud Dextrose Broth

LQ129V

Intended use

Recommended for cultivation of yeasts, moulds and aciduric bacteria.

Composition**

Ingredients	g / L
Peptone, special	10.000
Dextrose (Glucose)	20.000
Final pH (at 25°C)	5.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Label the ready to use LQ129V bottle. Inoculate the sample and Incubate at specified temperature and time.

Principle And Interpretation

Sabouraud Dextrose Agar is Carliers modifications (1) of the formulation described by Sabouraud (2) for the cultivation of fungi, particularly those associated with skin infections. The medium is also recommended by APHA (3). Sabouraud Dextrose Broth is also a modification by Sabouraud (4) and serves the same purpose as Sabouraud Dextrose Agar Medium 3.

Sabouraud dextrose media are peptone media supplemented with dextrose to support the growth of fungi. Peptone special provides nitrogen, vitamins, minerals, amino acids and growth factors. Dextrose (Glucose) provides an energy source for the growth of microorganisms. The low pH favors fungal growth and inhibits contaminating bacteria from clinical specimens (5). The acid reaction of the final medium is inhibitory to a large number of bacteria making it particularly useful for cultivating fungi and aciduric microorganisms. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously. Incubate cultures for 4 to 6 weeks before reporting as negative.

Type of specimen

Clinical samples: skin lesions, oral lesions, oropharyngeal swabs, vaginal secretions, urine, skin and nail scrapings, urine etc.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (6,7).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use only. For professional use only. Read the label before opening the container. Wear protective gloves/ protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Some fungi may show poor growth due to nutritional variations.
2. Further isolation and biochemical tests should be carried out for confirmation.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Sterile clear Sabouraud Dextrose Broth in bottle.

Colour

Light amber coloured clear solution.

Quantity of Medium

5ml of medium in glass bottle.

pH

5.40- 5.80

Sterility Check

Passes release criteria

Growth Promotion Test

Cultural characteristics was observed after an incubation at 30-35°C for 3-5 days.

Organism	Inoculum (CFU)	Growth
<i>Candida albicans</i> ATCC 10231 (00054*)	50-100	luxuriant
<i>Candida albicans</i> ATCC 2091 (00055*)	50-100	luxuriant
# <i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	50-100	luxuriant
<i>Saccharomyces cerevisiae</i> ATCC 9763 (00058*)	50-100	luxuriant
^ <i>Saccharomyces kudriavzevii</i> ATCC 2601	50-100	good-luxuriant
<i>Trichophyton rubrum</i> ATCC 28191	50-100	luxuriant (for 5-7 days)

Key : (*) Corresponding WDCM numbers,

(#) Formerly known as *Aspergillus niger*

(^) Formerly known as *Saccharomyces cerevisiae*

Storage and Shelf Life

On receipt store between 15-30°C. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (6,7).

Reference

- 1.Carlier G. I. M., 1984, Brit. J. Derm. Syph., 60:61
- 2.Sabouraud R., Les Teignes, Paris: Masson et Cie, 1910, p 553
- 3.Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
- 4.Sabouraud R., 1892, Ann. Dermatol. Syphil. 3 : 1061.
- 5.Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
- 6.Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
- 7.Jorgensen,J.H., Pfaller , M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.


Revision : 02/2024




HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India



**In vitro diagnostic
medical device**



Storage temperature



CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



CE Marking



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.