

Medium 11.GN Broth

LQ151C

Intended use

Recommended for the enrichment of *Shigella* from pharmaceutical products in accordance with IP 2022.

Composition**

Ingredients	g/ L
Polypeptone peptone	20.000
Glucose (Dextrose)	1.000
Sodium citrate	2.000
Sodium deoxycholate	0.500
Di-potassium hydrogen phosphate	4.000
Potassium dihydrogen phosphate	1.500
Sodium chloride	5.000
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Label the ready to use LQ151C bottle. Inoculate the sample and Incubate at specified temperature and time.

Principle And Interpretation

GN Broth is recommended by the Indian Pharmacopoeia (1) for the selective enrichment of *Shigella* species with subsequent isolation on a selective medium, XLD Agar (MH031). Croft and Miller isolated more strains of *Shigella* from rectal swabs using this medium (2). Hajna (3,4) also suggested the enrichment of organisms from rectal swabs in this medium 1-6 hours before plating on solid media.

The medium contains polypeptone peptone, which provides amino acids and other nitrogenous substances to support bacterial growth. The combination of sodium citrate and sodium deoxycholate inhibit gram-positive and some gram-negative bacteria such as coliforms. Phosphates serve as a buffering system. Sodium chloride maintains osmotic equilibrium. *Proteus*, *Pseudomonas* and coliforms do not overgrow *Salmonella* and *Shigella* in GN Broth during the first 6 hours of incubation. This enrichment broth should be used in conjunction with selective and non selective plating media to increase the probability of isolating pathogens (5,6,7).

Type of specimen

Pharmaceutical samples

Specimen Collection and Handling

For pharmaceutical samples follow appropriate techniques for handling specimens as per established guidelines (1).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Further biochemical and serological tests must be carried out for complete identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Sterile clear GN Broth in glass bottle.

Colour

Light amber coloured, clear to slightly opalescent solution.

Quantity of medium

100 ml of medium in glass bottles.

pH

6.80-7.20

Sterility Check

Passes release criteria

Growth promoting properties

Growth Promotion was carried out in accordance with the method of IP. Clearly visible growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≤ 100 cfu (at 30-35°C for 18 hours.)

Inhibitory Properties

No growth of the test microorganism occurs for the specified temperature for not less than longest period of time specified inoculating ≥ 100 cfu(at least 100 cfu)(at 30-35°C for ≥ 24 hours).

Organism	Inoculum (CFU)	Growth in GN broth	Recovery on XLD Agar	Recovery	Colour of colony
Growth promoting					
<i>Shigella boydii</i> ATCC 8700	50 -100	good	good-luxuriant	≥ 50 %	red translucent
Inhibitory					
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	$\geq 10^3$	inhibited		0 %	

Key (*) - Corresponding WDCM number

Storage and Shelf Lif

On receipt store between 15-30°C. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (8,9).

Reference

- 1.Indian Pharmacopoeia, 2022, Amd.2024 Indian Pharmacopoeia Commission, Ministry of Health and Family Welfare Government of India.
- 2.Croft C. C., Miller M. J., 1956, Am. J. Clin. Pathol., 26:411.
- 3.Hajna A. A., 1955, Publ. Health Lab., 13:83.
- 4.Hajna A. A., 1956, Air. Univ. Sch. Ar. Med., USAF, 56:39.
- 5.MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
- 6.Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
- 7.Salfinger Y., and Tortorello M.L. 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
- 8.Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
- 9.Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 06/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.