



Technical Data

MacConkey Agar w/o CV, NaCl, w/ 0.5% Sodium Taurocholate M082

Intended Use:

Recommended for the cultivation and differentiation of enteric bacteria, restricting swarming of *Proteus* species from specimens such as urine which may contain large number of *Proteus* species as well as potentially pathogenic Gram-positive organisms.

Composition**

Ingredients	g / L
Peptone	20.000
Lactose	10.000
Sodium taurocholate	5.000
Neutral red	0.040
Agar	20.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 55.0 grams of medium in 1000 ml purified/distilled water. Heat to boiling with gentle swirling to dissolve the agar completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Avoid overheating. Cool to 45-50°C. Mix well and pour into sterile Petri plates. The surface of the medium should be dry when inoculated.

Principle And Interpretation

MacConkey Agar is the earliest selective and differential medium for cultivation of enteric microorganisms from a variety of clinical specimens (1,2). Subsequently MacConkey Agar and Broth have been recommended for use in microbiological examination of foodstuffs (3) and for direct plating / inoculation of water samples for coliform counts (4). These media are also accepted by the Standard Methods for the Examination of Milk and Dairy Products (5) and pharmaceutical preparations (6).

Original medium contains protein, bile salts, sodium chloride and two dyes. The selective action of this medium is attributed to bile salts, which are inhibitory to most species of gram-positive bacteria. MacConkey Agar w/o CV, NaCl and W/ 0.5% Sodium taurocholate is a modification of the original formulation with the exclusion of crystal violet and inclusion of sodium taurocholate instead of bile salts. Gram-negative bacteria usually grow well on the medium and are differentiated by their ability to ferment lactose. Lactose fermenting strains grow as red or pink and may be surrounded by a zone of acid precipitated bile. The red colour is due to production of acid from lactose, absorption of neutral red and a subsequent colour change of the dye when the pH of medium falls below 6.8. Lactose non-fermenting strains, such as *Shigella* and *Salmonella* are colourless and transparent and typically do not alter appearance of the medium. *Yersinia enterocolitica* may appear as small, non-lactose fermenting colonies after incubation at room temperature.

Type of specimen

Clinical samples - faeces, urine, Food and Dairy samples, Water samples.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (7,8).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (3,5).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards (4). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/pr protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. The medium differentiates organisms on the basis of lactose fermentation. Further biochemical test must be carried out for confirmation.
2. The surface of the medium should be dry before inoculation.
3. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
4. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 2.0% Agar gel.

Colour and Clarity of prepared medium

Orange red coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 5.5% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony
<i>Salmonella</i> Paratyphi B ATCC 8759	50-100	luxuriant	≥50%	colourless
<i>Salmonella</i> Typhi ATCC 6539	50-100	luxuriant	≥50%	colourless
<i>Salmonella</i> Enteritidis ATCC 13076 (00030*)	50-100	luxuriant	≥50%	colourless
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	fair-good	30-40%	pale pink -red
<i>Salmonella</i> Paratyphi A ATCC 9150	50-100	luxuriant	≥50%	colourless
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	luxuriant	≥50%	pink to red with bile precipitate
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	50-100	fair to good	30-40%	pale pink to red
<i>Shigella flexneri</i> ATCC 12022 (00126*)	50-100	fair to good	30-40%	colourless
# <i>Klebsiella aerogenes</i> ATCC 13048 (00175*)	50-100	luxuriant	≥50%	pale pink to red
<i>Proteus vulgaris</i> ATCC 13315	50-100	luxuriant	≥50%	colourless

Key : *Corresponding WDCM numbers. #- Formerly known as *Enterobacter aerogenes*

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

Reference

1. MacConkey, 1900, The Lancet, ii:20.
2. MacConkey, 1905, J. Hyg., 5:333.
3. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
4. Lipps WC, Braun-Howland EB, Baxter TE, eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.
5. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
6. The United States Pharmacopoeia-National Formulary (USP-NF), 2022.
7. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
8. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 06/2024



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India




CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



In vitro diagnostic
medical device



CE Marking



Storage temperature



Do not use if
package is damaged

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.