

Technical Data

Fungal Agar (Mycological Agar)

M094

Intended Use:

For cultivation and maintenance of fungi.

Composition**

Ingredients	g/L
Soya peptone	10.000
Dextrose (Glucose)	10.000
Agar	15.000
Final pH (at 25°C)	7.0±0.2

^{**}Formula adjusted, standardized to suit performance parameters

Directions

Suspend 35 grams in 1000 ml purified / distilled water. Heat, to boiling, to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates. For preparing selective media, acidify the media upto pH 3.0-4.0 by the addition of two vials of 10% Lactic Acid Solution (FD095).

Principle And Interpretation

Mycological media are basal media to which antifungal agents may be added for checking their effect on fungi or bacteria to render them selective for isolation and cultivation of fungi. Mycological Agar is used while working with pathogenic fungi. Earlier media for fungi generally relied on an acidic pH to make the media less suitable for the growth of many bacteria (1). Fungal Agar is prepared according to the formulation suggested by Huppert and Walker (2).

Soya peptone in the medium provides nitrogen, vitamins and minerals necessary to support bacterial growth. Dextrose is a carbon source required for the growth of fungi. The pH may be adjusted to 4.0 after autoclaving by adding sterile 10% lactic acid sodium (FD195)/acetic acid and used for determining yeast and mould counts of carbonated beverages and food products (3).

Type of specimen

Clinical samples - nail scrappings, skin scrapings, etc.; Food samples.

Specimen Collection and Handling:

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (4,5). For food samples, follow appropriate techniques for sample collection and processing as per guidelines (6).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions:

In Vitro diagnostic Use. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations:

- 1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
- 2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

HiMedia Laboratories Technical Data

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.5% w/v aqueous solution at 25°C. pH: 7.0±0.2

pН

6.80-7.20

Cultural Response

Cultural characteristics observed after an incubation at 25 - 30°C for 48 - 72 hours (For *Trichophyton* species longer incubation may be required for upto 7days).

Organism	Inoculum (CFU)	Growth	Recovery
Growth Promotion # Aspergillus brasiliensis ATCC 16404 (00053*)	50-100	luxuriant	
Candida albicans ATCC 10231 (00054*)	50-100	luxuriant	>=70%
Lactobacillus acidophilus ATCC 11506	50-100	luxuriant	>=70%
Saccharomyces cerevisiae ATCC 9763 (00058*)	50-100	luxuriant	>=70%
Saccharomyces uvarum ATCC 28098	50-100	luxuriant	>=70%
Staphylococcus aureus subsp. aureus ATCC 25923 (00034*)	50-100	luxuriant	>=70%
Trichophyton mentagrophytes ATCC 9533	50-100	luxuriant	

Key: *Corresponding WDCM numbers. # - Formerly known as Aspergillus niger

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference

- 1. A. J. Clin. Path., 1951, 21: 684.
- 2. Huppert M., and Walker L. J., 1958, Am. J. Clin. Pathol., 29:291
- 3. Speck M. L., (Eds.), 1984, Compendium of Methods for the Microbiological Examination of Foods, 2nd Ed., APHA, Washington, D.C.
- 4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
- 5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
- 6. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.

HiMedia Laboratories Technical Data

Revision: 05/2024



HiMedia Laboratories Pvt. Limited, Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) -400604, MS, India



CEpartner4U, Esdoornlaan 13, 3951DB Maarn, NL www.cepartner4u.eu



In vitro diagnostic medical device





Storage temperature



Do not use if package is damaged

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.