

Technical Data

ITC Broth Base (TTC Broth Base)

Intended Use:

Recommended for selective enrichment and enumeration of *Yersinia enterocolitica*. The composition and performance criteria of this medium are as per the specifications laid down in ISO 10273:2017 and 11133:2014 (E) /Amd. :2020.

Composition**

ISO Specification - Irgasan, ticarcillin and potassium chlorate (ITC) broth		M1220 - ITC Broth Base (TTC Broth Base)		
Ingredients	g / L	Ingredients	g / L	
Enzymatic digest of casein	10.000	Tryptone #	10.000	
Yeast extract	1.000	Yeast extract	1.000	
Magnesium chloride hexahydrat	e 60.000	Magnesium chloride hexahydrate	60.000	
Sodium chloride	5.000	Sodium chloride	5.000	
Malachite green	0.010	Malachite green	0.010	
Triclosan (Irgasan)	0.001	Triclosan (Irgasan)	0.001	
Final pH (at 25°C)	6.9±0.2	Final pH (at 25°C)	6.9±0.2	
		Tic Selective Supplement (FD10)	2)- 1 vial	
Ticarcillin	1 mg	Ticarcillin	1 mg	
		PC Selective Supplement (FD10	3) - 1 vial	
Potassium chlorate	1 g	Potassium chlorate Distilled water	1 g 10.00 ml	

**Formula adjusted, standardized to suit performance parameters

Equivalent to Enzymatic digest of casein

Directions

Suspend 44.12 gram (the equivalent weight of dehydrated medium) in 988 ml purified/distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs (121°) for 15 minutes. Cool to 45-50°C. Aseptically add rehydrated contents of one vial of Tic Selective Supplement (FD102) and PC Selective Supplement (FD103). Mix well and dispense into sterile tubes or flasks as desired.

Principle And Interpretation

The genus *Yersinia* belongs to the family *Enterobacteriaceae*. They are usually nitrate reductase positive and show fermentative metabolism. The genus comprises of 11 species, of which *Yersinia enterocolitica* is most important as a causative agent of human foodborne illness. Variety of enrichment methods has been described for recovery of *Y.enterocolitica* from foods. The most efficient procedures for recovering enteropathogenic bacteria from foods have incorporated at least one and often two enrichment steps before plating onto selective differential agar media. ITC Broth is formulated in accordance with APHA (1) and is recommended by ISO Committee (2,3) as a selective enrichment medium for *Y.enterocolitica* from foods. ITC Broth was developed by Wauters et al (4) as a new enrichment broth, derived from modified Rappaport Broth and based on the selective agents irgasan (ticarcillin) and potassium chlorate.

Tryptone and yeast extract provide nitrogeneous and carbonaceous compounds, long chain amino acids and other essential growth nutrients. Ticarcillin has inhibitory action on both gram-positive and gram-negative organisms. Irgasan inhibits gram-positive organisms. Potassium chlorate has disinfecting properties.

Type of specimen

Food and animal feeding stuff

Specimen Collection and Handling:

Processing (1-3)

Enrichment : For the first initial suspension place the sample (x) in known volume of the PSB broth (M941I), to give a dilution of 1/10 dilution (by mass/volume or volume/volume). Homogenize the suspension using a peristaltic blender for 2 min. Incubate at 22°C to 25°C for 2 to 3 days with or 5 days without agitation.

M1220

For the second initial suspension in the same way with the ITC broth (M1220) so as to obtain a test portion/enrichment medium dilution of 1/100 (mass/volume or volume/volume). Incubate at 25°C for 48 hours. **Isolation** :

1.Inoculate the culture obtain from PSB culture on the surface of CIN agar plate and incubate at 30°C for 24 to 48 hours. 2.Alkaline treatment : Using sterile pipette transfer 0.5ml of the PSB culture into 4.5 ml of KOH solution and mix for 20 seconds only. Immediately inoculate on CIN agar plate. Incubate at 30°C for 24 to 48 hours.3.Using ITC culture inoculate the surface of CIN agar plate (M843I). Incubate at 30°C for 24 to 48 hours.

Purification : Streak the selected colonies on the surface of Nutrient Agar (M561A). Incubate at 30°C for 24 hours. Confirmation : Biochemical and serological tests are performed for confirmation.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to light blue homogeneous free flowing powder

Colour and Clarity of prepared medium

Peacock green coloured, clear solution without any precipitate

Reaction

Reaction of 4.41% w/v aqueous solution at 25°C. pH : 6.9±0.2

pН

6.70-7.10

Cultural Response

Productivity : Cultural characteristics observed with added Tic Supplement (FD102) and PC Supplement (FD103) after an incubation at 25±1°C for 44±4 hours. Recovery is carried out on CIN Agar (M843I) and characteristic reaction noted. **Selectivity:** Cultural characteristics observed with added Tic Supplement (FD102) and PC Supplement (FD103) after an incubation at 25±1°C for 44±4 hours. Recovery is carried out on Soyabean Casein Digest Agar (M290).

Organism	Inoculum (CFU)	Growth	Characteristic reaction of target organism on CIN Agar (M8431)
Productivity			5 5 , ,
<i>Yersinia enterocolitica</i> subsp. <i>palearctica</i> serotype O:3	50-100	>10 colonies	Transparent or translucent circular, smooth colonies with deep red sharp bordered centre.
NCTC 13769 (00126)*+ Escherichia coli ATCC 25922 (00013*)	>=10 ⁴		
Pseudomonas aeruginosa ATCC 27853 (00025*)	>=10 ⁴		
<i>Yersinia enterocolitica</i> subsp. <i>palearctica</i> serotype	50-100	>10 colonies	Transparent or translucent circular, smooth colonies with deep red sharp bordered centre.
O:3 NCTC 13769 (00126)* + <i>Escherichia coli</i> ATCC	>=10 ⁴		
8739 (00012*) + <i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	>=10 ⁴		
<i>Yersinia enterocolitica</i> ATCC 23715 (00160)*	50-100	>10 colonies	Transparent or translucent circular, smooth colonies with deep red sharp bordered centre.
+ <i>Escherichia coli</i> ATCC 25922 (00013*)	>=10 ⁴		
+ Pseudomonas aeruginosa ATCC 27853 (00025*)	>=10 ⁴		

Yersinia enterocolitica ATCC 23715 (00160)* + Escherichia coli ATCC 8739 (00012*)	50-100	>10 colonies	Transparent or translucent circular, smooth colonies with deep red sharp bordered centre.
+ Pseudomonas aeruginosa > ATCC 27853 (00025*)	>=104		
Selectivity			
Pseudomonas aeruginosa	>=10 ⁴	inhibited	
ATCC 27853 (00025*)			
Proteus mirabilis ATCC	>=10 ⁴	inhibited	
29906 (00023*)			

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

1. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.

2. ISO 10273:2017 Microbiology of the food chain Horizontal method for the detection of pathogenic Yersinia enterocolitica 3. Microbiology of food, animal feeding stuffs and water- Preparation, production, storage and performance testing of culture media, EN ISO 11133:2014 (E) /Amd.2 :2020.

4. Wauters G., Goossens V., Janssens M. and Vandepitte J., 1988, In. J. Syst. Bacteriol., 38, 424-429.

5. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.

6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 08/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.