



## Mucate Control Broth

M1227

Mucate Control Broth is used for identification of enteropathogenic *Escherichia coli* and *Salmonella* species from milk and milk products.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Bromothymol blue	0.024
Final pH ( at 25°C)	7.4±0.1

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 10 grams in 1000 ml distilled water. Dispense in 5 ml amounts in screw-capped tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes.

### Principle And Interpretation

Mucate Broth is prepared based on the formula originally developed by Kauffman and Petersen (1) recommended by APHA (2) for identification of enteropathogenic *Escherichia coli* from milk and milk products. This medium can also be used as an aid in differentiation of *Enterobacteriaceae* especially within *Salmonella genus* .

Mucic acid is a saccharolactic acid or also called as tetrahydroxyadipic acid and act as a carbon source in the medium. It is fermented by enteropathogenic *Escherichia coli* , *Salmonella Paratyphi B* and also by *Klebsiella pneumoniae* to produce acid which makes the medium yellow as the pH indicator is bromo thymol blue (3). Peptic digest of animal tissue supplies the necessary nutrients to the organisms.

Transfer a loopful of 24 hour Tryptone Broth (M463) culture to Mucate Broth. Include Mucate Control Broth tube as a control because occasionally un-inoculated tubes of Mucate Broth turn blue on standing. Incubate at  $48 \pm 1$  hour at 35°C. A negative test result is indicated by a blue or unchanged colour in this broth. 90 % of the *E.coli* strains are mucate positive.

### Quality Control

#### Appearance

Cream to pale green homogeneous free flowing powder

#### Colour and Clarity of prepared medium

Blue coloured clear solution without any precipitate

#### Reaction

Reaction of 1.0% w/v aqueous solution at 25°C. pH : 7.4±0.1

#### pH

7.30-7.50

#### Cultural Response

M1227: Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours on addition of 1% Mucic acid.

Organism	Inoculum (CFU)	Growth
<b>Cultural Response</b>		
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	luxuriant
<i>Salmonella Paratyphi B</i>	50-100	luxuriant

### Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Kauffmann F., and Petersen A., 1956, Acta. Pathol. Microbiol. Scand., 38 (6) : 481.
2. Marshall R. (Ed.), 1992, Standard Methods for the Examination of Dairy Products, 16th ed., APHA, Washington, D.C.
3. MacFaddin J.F., 1985, Media for Isolation - Cultivation - Identification - Maintenance of Medical Bacteria, Vol.I, Williams and Wilkins, Baltimore.

Revision : 1 / 2011



### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.