



HiCrome™ Coliform Agar w/ SLS

M1300

Intended use

Recommended for the simultaneous detection of *Escherichia coli* and total coliforms in water, food and clinical samples.

Composition**

Ingredients	g / L
Peptone, special	3.000
Sodium chloride	5.000
Dipotassium hydrogen phosphate	3.000
Potassium dihydrogen phosphate	1.700
Sodium pyruvate	1.000
L-Tryptophan	1.000
Sodium lauryl sulphate	0.100
Chromogenic mixture	0.200
Agar	12.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 27.0 gram in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Add 5mg/l novobiocin before autoclaving the medium, when a high number of gram positive accompanying bacteria are expected. Mix well and pour into sterile Petri plates.

Principle And Interpretation

HiCrome™ Coliform Agar w/ SLS is a selective medium recommended for the simultaneous detection of *Escherichia coli* and total coliforms in water and food samples (1). Peptone special and sodium pyruvate provide essential growth nutrients to the organisms. The phosphates buffer the medium well. The medium composition helps even the sub lethally injured coliforms to grow rapidly. Sodium lauryl sulphate inhibits gram-positive organisms. The chromogenic mixture contains two chromogenic substrates. The enzyme β -galactosidase produced by coliforms cleaves one chromogen, resulting in the salmon red colouration of coliform colonies. The enzyme β -glucuronidase produced by *E. coli*, cleaves X-glucuronide. *E.coli* forms dark blue to violet coloured colonies due to cleavage of both the chromogens (2,3,4). The addition of L-Tryptophan improves the indole reaction, thereby increasing detection reliability in combination with the two chromogens. To confirm *E.coli*, add a drop of Kovac's reagent (R008) on the dark-blue to violet colony. Formation of cherry-red colour indicates positive reaction.

Type of specimen

Clinical samples - faeces, urine, etc.; Food samples; Water samples.

Specimen Collection and Handling:

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (5,6).

For food samples, follow appropriate techniques for sample collection and processing as per guidelines (7).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards (8).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. β -glucuronidase is present in 97% of *E.coli* strains, however few *E.coli* may be negative.
2. Certain species of *Shigella* and *Salmonella* are β -glucuronidase positive.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to beige homogeneous free flowing powder

Gelling

Firm, comparable with 1.2% Agar gel.

Colour and Clarity of prepared medium

Light yellow, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 2.7% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH

6.60-7.00

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24 hours (48 hours if necessary).

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony	Indole production
<i>Citrobacter freundii</i> ATCC 8090	50-100	good-luxuriant	≥50%	salmon to red	negative reaction
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	good-luxuriant	≥50%	dark blue/violet	positive, confirmation of red colour around the colony by addition of Kovacs reagent (R008)
<i>Enterobacter cloacae</i> ATCC 23355 (00082*)	50-100	good-luxuriant	≥50%	salmon to red	negative reaction
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	≥10 ⁴	inhibited	0%		
<i>Klebsiella pneumoniae</i> ATCC 13883 (00097*)	50-100	good-luxuriant	≥50%	light pink	negative reaction
<i>Salmonella</i> Enteritidis ATCC 3076 (00030*)	50-100	good	40-50%	colourless	negative reaction
<i>Shigella flexneri</i> ATCC 12022 (00126*)	50-100	good	40-50%	colourless	negative reaction

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 15-25°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

1. Manafi M. and Kneifel W., 1989, Zentralbl. Hyg., 189:225.
2. Frampton E. W., Restaino L. and Blaszkowski N., 1988, J. Food Prot., 51:402.
3. Kilian M. and Bülow P., 1976, Acta. Pathol. Microbiol. Scand., Sect. B, 84:245.
4. LeMinor L. and Hamida F., 1962, Ann. Inst. Pasteur (Paris), 102:267
5. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
7. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
8. Lipps WC, Braun-Howland EB, Baxter TE, eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.C


Revision : 05/2024



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India



**In vitro diagnostic
medical device**



Storage temperature



CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



CE Marking



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.