



Sabouraud Dextrose Maltose Agar

M1313

Intended Use:

Recommended for cultivation of moulds and yeasts and for testing antimycotic substances.

Composition**

Ingredients	g / L
Tryptone	5.000
Peptone	5.000
Dextrose (Glucose)	10.000
Maltose	10.000
Agar	15.000
Final pH (at 25°C)	5.4±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 45.0 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. **DO NOT OVERHEAT**. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Sabouraud Dextrose Agar is Carliers modifications (1) of the formulation described by Sabouraud (2) for the cultivation of fungi, particularly those associated with skin infections. Sabouraud Dextrose Maltose Agar is used for the cultivation of yeast, moulds and other aciduric organisms (3,4,5).

Sabouraud dextrose media are peptone media supplemented with dextrose to support the growth of fungi. Tryptone and peptone provide nitrogen, vitamins, minerals, amino acids and growth factors. Dextrose and maltose provide an energy source for the growth of microorganisms. The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens (6). The acid reaction of the final medium is inhibitory to a large number of bacteria making it particularly useful for cultivating fungi and aciduric microorganisms. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously. Incubate cultures for 4 to 6 weeks before reporting as negative.

Type of specimen

Clinical samples - Skin scrapping or swab of infected region; Water samples

Specimen Collection and Handling:

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (7,8).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(9) After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/ protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.
3. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control**Appearance**

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.5% w/v aqueous solution at 25°C. pH : 5.4±0.2

pH

5.20-5.60

Cultural Response

Cultural characteristics observed after an incubation at 25 - 30°C for upto 5 days.

Organism	Inoculum (CFU)	Growth	Recovery
# <i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	50-100	good-luxuriant	
<i>Candida albicans</i> ATCC 10231 (00054*)	50-100	good-luxuriant	≥70%
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	good-luxuriant(Inhibited on media with low pH)	≥70%
<i>Lactobacillus casei</i> ATCC 9595	50-100	good-luxuriant	≥70%
<i>Saccharomyces cerevisiae</i> ATCC 9763 (00058*)	50-100	good-luxuriant	≥70%
<i>Trichophyton rubrum</i> ATCC 28191	50-100	good-luxuriant	
<i>Penicillium notatum</i> ATCC 10108	50-100	Good-luxuriant	
<i>Trichophyton gallinae</i> ATCC 22243	50-100	Good-luxuriant	
<i>Trichophyton mentagrophytes</i> ATCC 9533	50-100	Good-luxuriant	
<i>Trichophyton ajelloi</i> ATCC 24885	50-100	Good-luxuriant	

Key: *Corresponding WDCM numbers. #Formerly known as *Aspergillus niger*

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (3,4).

Reference

1. Carlier G. I. M., 1984, Brit. J. Derm. Syph., 60:61
2. Sabouraud R., 1892, Ann. Dermatol. Syphil. 3 : 1061.
3. Merkblatt 18: Verpackgs- Rdsch, 1974, 25/1: Techn- Wiss. Beilage, 5-8
4. Merkblatt 19: Verpackgs- Rdsch, 1974, 25/6: Techn- Wiss. Beilage, 569-575.

5. Merkblatt 21: Verpackgs- Rdsch, 1974, 25/7: Techn- Wiss. Beilage, 53-55
6. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenenbaum R. H., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
7. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
8. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
9. Lipps WC, Braun-Howland EB, Baxter TE, eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.


Revision : 03/2024



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India



**In vitro diagnostic
medical device**



Storage temperature



CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



CE Marking



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.