

# **Kings Medium A Base**

# Intended Use:

For non-selective isolation, cultivation and pigment production of Pseudomonas species.

Composi	ition	**
---------	-------	----

Ingredients	g / L
Proteose peptone	20.000
Potassium sulphate	10.000
Magnesium chloride	1.640
Agar	15.000
Final pH ( at 25°C)	7.3±0.1

\*\*Formula adjusted, standardized to suit performance parameters

## Directions

Suspend 46.64 grams in 1000 ml purified/distilled water containing 10 ml of glycerol. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and aseptically pour into sterile Petri plates.

# **Principle And Interpretation**

*Pseudomonas aeruginosa* is known to produce two types of pigments, pyocyanin and fluorescein which is a characteristic property and aids in isolation of *Pseudomonas* from clinical material. An additional pigment called as pyorubin was reported by King. Pyocyanin is green while fluorescein is fluorescent yellow and pyorubin is reddish brown. Some strains produce all these pigments while the others produce one or two pigments. *P.aeruginosa* can be identified on Hugh Leifson Medium (M826). Kings Medium A Base is particularly suited for the production of pyocyanin and pyorubin. Kings Medium A Base is based on the formulation of King et al (1,2). This medium can be used as a general medium for the non-selective isolation and pigment production of *Pseudomonas* species from foods, cosmetic samples etc. These media contain proteose peptone, which provides carbonaceous and nitrogenous compounds for the growth of bacteria. Glycerol serves as a source of energy and also enhances pigment production. Magnesium chloride, potassium sulphate and magnesium sulphate also enhances pigment production. Pigments and/ or their derivatives produced by *Pseudomonas* species play a role as siderophores in the iron uptake systems of bacteria, and hence, their production is markedly enhanced under conditions of iron deficiency. For inoculation, use the organisms freshly cultured in Kings Medium A, incubate overnight at 37°C and then at room temperature for 6 days.

## **Type of specimen**

Clinical samples - pus, urine ; Water samples.

# **Specimen Collection and Handling:**

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (3,4). For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards (5). After use, contaminated materials must be sterilized by autoclaving before discarding.

## **Warning and Precautions**

In Vitro diagnostic use. For professional use only. Read the label before opening the container. The media should be handled by trained personnel only. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

## **Limitations :**

1. For inoculation, use the organisms freshly cultured in Kings Medium A, incubate overnight at  $37^{\circ}$ C and then at room temperature for 6 days.

2. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.

## **Performance and Evaluation**

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## **Quality Control**

## Appearance

Cream to yellow homogeneous free flowing powder

## Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 4.6% w/v aqueous solution (containing 1.0%v/v glycerol) at 25°C. pH : 7.3±0.1

#### pН

7.20-7.40

#### **Cultural Response**

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Pigment production
Pseudomonas aeruginosa ATCC 17934	50-100	good-luxuriant	>=70%	blue-green
Pseudomonas aeruginosa ATCC 27853 (00025*)	50-100	good-luxuriant	>=70%	blue-green
Pseudomonas aeruginosa ATCC 9027 (00026*)	50-100	good-luxuriant	>=70%	blue-green
<i>Burkholderia cepacia</i> ATCC 25609	50-100	good-luxuriant	>=70%	no pigment

Key: (\*) - Corresponding WDCM numbers

## Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (3,4).

## Reference

- 1. King E. O., Ward M. K. and Raney D. E., 1954, J. Lab and Clin. Med., 44:301-307.
- 2. Murray P. R., Baron E. J., Jorgensen J. H., Pfaller M.A., Yolken R. H., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
- 3. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
- Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
- 5. Lipps WC, Braun-Howland EB, Baxter TE,eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.

Revision : 04/2024





HiMedia Laboratories Pvt. Limited, Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) -400604, MS, India

CEpartner4U, Esdoornlaan 13,

3951DB Maarn, NL

www.cepartner4u.eu



IVD



-30°C Storage temperature

> Do not use if package is damaged

#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia<sup>TM</sup> publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia<sup>TM</sup> Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

In vitro diagnostic

HiMedia Laboratories Pvt. Ltd. Corporate Office : Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) - 400604, India. Customer care No.: 022-6147 1919 Email: techhelp@himedialabs.com Website: www.himedialabs.com