



Letheen Broth I Modified

Intended Use:

Recommended for screening cosmetic products for microbial contamination.

Composition**

Ingredients	Gms / Litre
HM peptone B #	5.000
Proteose peptone	20.000
Polysorbate 80 (Tween 80)	5.000
Lecithin	0.700
Sodium chloride	5.000
Tryptone	5.000
Yeast extract	2.000
Sodium bisulfite	0.100
Final pH (at 25°C)	7.2 ± 0.2
**Formula adjusted standardized to suit performance perameters	

**Formula adjusted, standardized to suit performance parameters

#- Equivalent to Beef extract

Directions

Suspend 42.8 grams in 1000 ml purified / distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and dispense as desired.

Principle And Interpretation

In the early 40s, Weber and Black recommended the use of lecithin and polysorbates to neutralize the antimicrobial action of the quaternary ammonium compounds (8). In 1965, the methodology was accepted by AOAC for the antimicrobial assays and extended their use to all the cationic detergents. In 1978, the FDA incorporated it as pre-enrichment medium for every microbial examination of cosmetics.

Letheen Broth, Modified is prepared as per FDA (1) for screening cosmetic products for microbial contamination. There are great chances of altering the chemical composition of cosmetics by the metabolism of organism s thereby spoiling and causing harm to the users (2, 7, 9). Direct colony counts and enrichment culturing are the methods of choice for isolating microorganisms from cosmetic products. The word Letheen represents a combination of lecithin and polysorbate (tween 80).

Proteose peptone, HM peptone B,tryptone and yeast extract provide nitrogenous nutrients, carbon compounds and trace elements to the microorganisms. Incorporation of lecithin and polysorbate 80 to the medium enables the recovery of bacteria from materials containing residues of disinfectant compounds or preservatives used in cosmetics. Polysorbate 80 is added to nullify phenolic compounds, hexachlorophene, formalin and along with lecithin neutralizes ethyl alcohol (3). Lecithin also neutralizes quaternary ammonium compounds present in the cosmetics. Sodium chloride maintains the osmotic balance of the medium.

Type of specimen

Cosmetic samples.

Specimen Collection and Handling

For cosmetics samples follow appropriate techniques for handling specimens as per established guidelines (4). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

Please refer disclaimer Overleaf.

1. Further biochemical and serological test must be carried out for further identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Yellow coloured, clear solution in tubes

Reaction

Reaction of 4.28% w/v aqueous solution at 25°C. pH : 7.2±0.2

pН

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth
Escherichia coli ATCC 25922 (00013*)	50-100	luxuriant
Staphylococcus aureus subsp. aureus ATCC 25923 (00034*)	50-100	luxuriant

Key : (*) Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

- 1. Bacteriological Analytical Manual, 2017, Food and Drug Administration, AOAC International, Gaithersburg, MD, U.S.A.
- 2. Dunningan A. P., 1968, Drug Cosmet. Ind., 102:43.
- 3. Favero (Chm.), 1967, A State of the Art Report, Biological Contamination Control Committee, American Association for Contamination Control.
- 4. Hall and Hartnett, 1964, Public Health. Rep., 79:1021.
- 5. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
- 6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
- of Clinical Microbiology, 11th Edition. Vol. 1.
- 7. Smart R. and Spooner D. F., 1972, J. Soc. Cosmet. Chem., 23:721.
- 8. Weber and Black, 1948, Soap Sanitary Chem., 24:134-139
- 9. Wilson L. A. and Ahearn D. G., 1977, Am. J. Opthalmol., 84:112.

Disclaimer :

Revision : 01 / 2019

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia[™] publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia[™] Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.