



Nutrient Broth with 1% Peptone

M244

Intended Use

Recommended as a sterility testing medium for aerobes.

Composition**

Ingredients	Gms / Litre
Peptone	10.000
HM peptone B#	10.000
Sodium chloride	5.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Equivalent to Beef extract

Directions

Suspend 25 grams in 1000 ml purified / distilled water. Heat if necessary to dissolve the medium completely. Dispense in tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Nutrient Broth with 1% Peptone has almost double concentration of the nitrogen sources than that used in Nutrient Broth, making it more nutritive.

Peptone and HM peptone B provide the necessary carbon, nitrogen compounds, vitamins and also some trace ingredients to non-fastidious organisms like *Bacillus subtilis* and *Staphylococcus aureus* (3). Sodium chloride maintains osmotic equilibrium of the medium.

Nutrient Broth with 1% Peptone can be used as a sterility testing medium for aerobes against Nutrient Broth recommended for microbial limit tests as per standard pharmacopoeia (6). This broth can also be used as the suspending medium for cooked meat granules for the cultivation of anaerobic organisms. Nutrient Broth w/ 1% Peptone is a nutritionally rich medium that facilitates the growth of very low inocula, when with fastidious microorganisms.

Type of specimen

Food and dairy samples ; Water samples

Specimen Collection and Handling

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,7,8).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards(2).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets

Limitations

1. Isolation on selective media is required for further confirmation.
2. Biochemical and serological testing needs to be done for complete identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution in tubes

Reaction

Reaction of 2.5% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	luxuriant
# <i>Klebsiella aerogenes</i> ATCC 13048 (00175*)	50-100	luxuriant
<i>Klebsiella pneumoniae</i> ATCC 13883 (00097*)	50-100	luxuriant
<i>Salmonella</i> Typhimurium ATCC 14028 (00031*)	50-100	luxuriant
<i>Escherichia coli</i> ATCC 8739 (00012*)	50-100	luxuriant
<i>Escherichia coli</i> NCTC 9002	50-100	luxuriant
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	50-100	luxuriant
<i>Salmonella</i> Abony NCTC 6017 (00029*)	50-100	luxuriant

Key : (*) Corresponding WDCM numbers. (#) Formerly known as *Enterobacter aerogenes*

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference

1. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
2. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
3. IP: Indian Pharmacopoeia, 2018, Govt. of India, 2018. The Controller of Publication, Delhi.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.

5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
6. MacFaddin J. F., 2000, Biochemical Tests for Identification of Medical Bacteria, 3rd Ed., Lippincott, Williams and Wilkins, Baltimore.
7. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
8. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.

Revision : 01 / 2019

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.