

Technical Data

Mycoplasma Agar Base (PPLO Agar Base)

Intended Use:

With the addition of enrichment it is recommended for isolation and cultivation of *Mycoplasma* species (Pleuro pneumonia- like organisms- PPLO).

Composition**

Ingredients	g / L
HM infusion B from 250 g #	6.000
Peptone	10.000
Sodium chloride	5.000
Agar	15.000
Final pH (at 25°C)	7.8±0.2

**Formula adjusted, standardized to suit performance parameters

Equivalent to Beef heart, infusion from

Directions

Suspend 36.0 grams in 700 ml purified / distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add 300 ml Horse serum (RM1239) or 10 vials of HPYT Enrichment Supplement (FD075). Mix well and pour into sterile Petri plates. 25% Ascitic fluid can be used instead of Horse serum.

Principle And Interpretation

PPLO Agar was described by Morton, Smith and Leberman (1). It was used in a study of the growth requirements of *Mycoplasma* (2), along with the identification and cultivation of this organism (3,4,5). Pivotal information regarding *Mycoplasma* has been documented by Sabin (6). Hayflick et al have reported the information regarding the cultivation of *Mycoplasma* (7).

For the cultivation of *Mycoplasma* the medium ingredients and all the supplements should be free of any toxic substances even in small amounts. HM infusion B from and peptone provide nitrogen and carbon source, long chain amino acids, vitamins, and other essential nutrients. Sodium chloride maintains the osmotic balance of these formulations. Many *Mycoplasma* require serum for their good growth and also presence of antibiotic is necessary to prevent the growth of contaminating organisms. Mostly the *Mycoplasma* species are aerobic or facultatively anaerobic but some are microaerophilic. Few are anaerobic saprophytic *Mycoplasma* which grow best at 22-35°C while pathogenic strains grow at 35°C. *Mycoplasma* when grow in the agar medium show typical morphology and form colonies below the agar surface and do no grow without serum.

Plates or tubes should be incubated in an atmosphere containing 5-10% carbon dioxide and examined after incubation of 48 hours but they should not be discarded as negative until after incubation for 3 weeks.

PPLO colonies are round with a dense center and a less dense periphery, resembling a "fried egg" on PPLO Agar. Vacuoles, large bodies characteristic of *Mycoplasma* species are seen in the periphery. Colonies vary in diameter from 10 to 500 microns (0.01-0.5 mm) and penetrate into the medium.

Type of specimen

Clinical samples- Urine sample, urethral swab, sputum sample; pharmaceutical samples

Specimen Collection and Handling:

For pharmaceutical samples, follow appropriate techniques for sample collection, processing as per pharmaceutical guidelines (8).

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (9,10). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use. For professional use only. Read the label before opening the container. Wear protective gloves/ protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and

M266

culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Since *Mycoplasma* species are aerobic or facultatively anaerobic but some are microaerophilic, proper incubation should be carried out for optimal recovery.

2.Few are anaerobic saprophytic Mycoplasma which grow best at 22-35°C while pathogenic strains grow at 35°C, hence growth conditions must be maintained.

3.Since the medium is highly enriched care must be taken during innoculation to avoid contamination.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.6% w/v aqueous solution at 25°C. pH : 7.8±0.2

pН

7.60-8.00

Cultural Response

Cultural characteristics observed in presence of 10% Carbon dioxide with added,1% Horse serum (RM1239) or 10 vials of HPYT Enrichment Supplement (FD075), after an incubation at 22-35°C for 48 hours.

Organism	Growth
<i>Mycoplasma bovis</i> ATCC 25523	good-luxuriant
<i>Mycoplasma gallinarium</i> ATCC 19708	good-luxuriant
<i>Mycoplasma pneumoniae</i> ATCC 15531	good-luxuriant
Streptococcus pneumoniae ATCC 6303	good-luxuriant

Storage and Shelf Life

Store below 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (9,10).

Reference

Morton, Smith and Leberman, 1951, Am. J. Syphilis Gonorrh. Veneral Diseases, 35: 361.
Morton and Lecce, 1953. J. Bacteriol., 66:646.
Chanock, James, Fox, Turner, Mufso and Hayflick, 1962, Soc. Exp. Biol. Med., 110:884.
Craven, Wenzel, Calhoun, Hendley, Hamory and Gwaltney, 1976, J. Clin. Microbiol., 4:225.
Gregory and Cundy, 1970, Appl. Microbiol., 19:268.

- 6. Sabin, 1941, Bacteriol. Rev., 5:1, 331.
- 7. Hayflick and Chanock, 1965, Bacteriol, Rev., 29:185.
- 8. The United States Pharmacopoeia-National Formulatory (USP-NF), 2022.
- 9. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
- 10. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 05/2024



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

HiMedia Laboratories Pvt. Ltd. Corporate Office : Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) - 400604, India. Customer care No.: 022-6147 1919 Email: techhelp@himedialabs.com Website: www.himedialabs.com