



Phenylethyl Alcohol Agar

M269

Intended Use:

For the selective isolation of gram positive organisms like *Staphylococci* and *Streptococci*.

Composition**

Ingredients	g / L
Tryptone	15.000
Soya peptone	5.000
Sodium chloride	5.000
Phenyl ethanol	2.500
Agar	15.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 42.5 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For the preparation of blood agar aseptically add 5% v/v sterile defibrinated blood to the sterile molten medium cooled to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Phenylethyl alcohol is a chemical agent that exhibits inhibitory action against gram-negative and certain gram-positive bacteria. Phenylethyl Alcohol Agar is formulated as per Lilley and Brewer (1) for the selective isolation of gram-positive bacteria. This medium can be supplemented with 5 % sheep blood. This medium is especially useful when specimens are contaminated with swarming *Proteus* species. It is also useful in the diagnostic studies of wounds and exudate cultures (2). However, Phenylethyl Alcohol Agar cant be used to study haemolytic reactions as the results are atypical.

Tryptone and soya peptone provide nitrogen, carbon, sulfur and trace elements to the growing organisms. Addition of sheep blood provides many growth factors. Sodium chloride maintains osmotic equilibrium. Addition of phenylethanol to a nutritive medium permits the growth of gram-positive organisms but inhibits the gram-negative organisms found in the same specimen (3). Phenylethyl alcohol exerts inhibitory bacteriostatic action on gram-negative bacteria by inhibiting their DNA synthesis (3).

Type of specimen

Clinical samples : pus and wounds swabs, faeces, etc.; Food samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (4,5).

For food samples, follow appropriate techniques for sample collection and processing as per guidelines (6).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Due to nutritional variations, some strains may show poor growth.
2. Phenylethyl Alcohol Agar cant be used to study haemolytic reactions as the results are atypical.
3. Further biochemical and serological tests must be performed for confirmation.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Basal medium: Light amber coloured clear to slightly opalescent gel. After addition of 5%v/v sterile defibrinated blood : Cherry red coloured opaque gel forms in Petri plates

Reaction

Reaction of 4.25% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH

7.10-7.50

Cultural Response

Cultural characteristics observed with added 5% v/v sterile defibrinated blood, after an incubation at 35-37°C for 18-48 hours

Organism	Inoculum (CFU)	Growth	Colour of colony
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	good-luxuriant	white to gray or cream to yellow
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	50-100	good-luxuriant	blue-gray
<i>Salmonella</i> Typhi ATCC 6539	50-100	none-poor	
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	none-poor	
<i>Proteus mirabilis</i> ATCC 25933	50-100	none-poor	

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store dehydrated and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (3,4).

Reference

1. Lilley B. D. and Brewer J. H., 1953, J. Am. Pharm. Assoc., 42:6.
2. Holzman J. A., 1958, Am. J. Med. Technol., 24 (5), 327,342.
3. Dowell, Hill and Altemeier, 1964, J. Bacteriol., 88:1811.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
6. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological of Examination Foods, 5th Ed., American Public Health Association, Washington, D.C.

Revision : 05/2024



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India




CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



**In vitro diagnostic
medical device**



CE Marking



Storage temperature



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.