

Technical Data

C.L.E.D. Agar w/ Andrade Indicator

Intended Use:

Recommended for isolation and differentiation of urinary pathogens on the basis of lactose fermentation. **Composition****

Ingredients	g / L
Peptone	4.000
HM peptone B #	3.000
Tryptone	4.000
Lactose	10.000
L-Cystine	0.128
Bromothymol blue	0.020
Andrade indicator	0.100
Agar	15.000
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

- Equivalent to Beef extract

Directions

Suspend 36.25 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Sandys reported a new technique where the swarming of *Proteus* on an agar medium could be prevented by restricting the electrolyte content in the culture medium (1). Sandys Medium was modified by Mackey and Sandys (2), by replacing mannitol with lactose and sucrose and elevating the concentration of agar and bromothymol blue. The same authors further modified this medium by retaining the lactose (deleting sucrose) and by including L-cystine for promoting the growth of cystine-dependent dwarf coliform colony (3). This later modified medium was designated as C.L.E.D. (Cystine- Lactose-Electrolyte-Deficient) Medium. This medium is recommended for use in urinary bacteriology, promoting the growth of all urinary pathogens. C.L.E.D. Medium is also recommended for dip stick procedures and as dip inoculum transport medium for urine specimens (2,3,4).

C.L.E.D. Medium was further modified by Bevis (5) by incorporation of Andrades indicator. This medium provides sharper differentiation between lactose-fermenters (LF) and lactose-non-fermenters (NLF) (5). Addition of Andrades indicator enhances the appearance of colony and aids in the identification of microorganisms.

At different pH values, the colour of the medium varies from the standard medium, which is well documented by Bevis (5).

pН	Colour of C.L.E.D. medium
1	

7.4	deep blue
	1

- 7.0 bluish grey
- 6.8 pale grey
- 6.6 pinkish grey
- 6.4 bright red with whitish tinge
- 6.0 bright red

For better results, the medium should not be incubated for more than 24 hours because if lactose fermenters predominate, the entire medium may turn pink masking the presence of non-lactose fermenters. Inoculate the medium immediately after urine collection. Shigella species may not grow on this medium. Prior initiation of antibiotic therapy, low urine pH (less than 5) etc. may result in low urine count from infected patients.

Type of specimen

Clinical:Urine sample

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (6,7). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use only. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. This medium is recommended for urine infection. Low urine count may be a result of antibiotic therapy, low pH of urine.

- 2. Recovery depends on the urine count.
- 3. Inoculate the medium immediately after urine collection.
- 4. Shigella species may not grow on this medium.

5. For better results, the medium should not be incubated for more than 24 hours because if lactose fermenters predominate, the entire medium may turn pink masking the presence of non-lactose fermenters.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to gravish yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Greenish blue clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.62% w/v aqueous solution at 25°C. pH : 7.5±0.2

pН

7.30-7.70

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
# Klebsiella aerogenes ATCC 13048 (00175*)	50-100	good-luxuriant	>=70%	greyish green
Escherichia coli ATCC 25922 (00013*)	50-100	good-luxuriant	>=70%	bright pink with pink halo
Enterococcus faecalis ATCC 29212 (00087*)	50-100	good-luxuriant	>=70%	orange-yellow or greenish
Proteus mirabilis ATCC 25933	50-100	good-luxuriant	>=70%	blue-green
Staphylococcus aureus subsp. aureus ATCC 25923 (00034*)	50-100	good-luxuriant	>=70%	golden-yellow
Streptococcus pyogenes	50-100	good-luxuriant	>=70%	greyish green

Key : *Corresponding WDCM numbers.

#- Formerly known as Enterobacter aerogenes

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (6,7).

Reference

- 1. Sandys, 1960, J. Med. Lab. Technol., 17:224.
- 2. Mackey and Sandys, 1965, Br. Med. J., 2:1286.
- 3. Mackey and Sandys, 1966, Br. Med. J., 1:1173.
- 4. Dixson J. M. S. and Clark M. A., 1968, Conc. Med. Assoc. J., 99 (15)
- 5. Bevis T. D., 1968, J. Med. Lab. Technol., 25:38.
- 6. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.

7. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 06/ 2024



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

HiMedia Laboratories Pvt. Ltd. Corporate Office : Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) - 400604, India. Customer care No.: 022-6147 1919 Email: techhelp@himedialabs.com Website: www.himedialabs.com