



Eugonic Broth

M429

Intended Use:

Recommended for cultivation of fastidious microorganisms like *Haemophilus*, *Neisseria*, *Pasteurella*, *Brucella* and *Lactobacillus* species.

Composition**

Ingredients	g / L
Tryptone	15.000
Soya peptone	5.000
Dextrose (Glucose)	5.000
Sodium chloride	4.000
Sodium sulphite	0.200
L-Cystine	0.200
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Equivalent to Papaic digest of soyabean meal

Directions

Suspend 29.4 grams in 1000 ml purified / distilled water. Heat if necessary with frequent stirring to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 40-45°C and add 5 -10% v/v sterile defibrinated blood if desired. The blood may be chocolate by heating.

Principle And Interpretation

Eugonic Broth was developed by Pelczar and Vera (1) for cultivation of fastidious organisms like *Brucella*. This medium can also be used to grow *Mycobacteria* and various pathogenic fungi including *Nocardia*, *Histoplasma* and *Blastomyces*, when enriched with blood. Niven used this media for detection of spoilage of meats (2). Eugonic Broth was developed to obtain eugonic (luxuriant) growth of fastidious microorganisms like *Brucella* which are otherwise difficult to cultivate (3). The unenriched medium supports rapid growth of lactobacilli associated with cured meat products, dairy products and other foods. APHA recommends Eugonic Broth, which is also used in germinating anaerobic spores pasteurized at 104°C (4,5). Organisms like *Bordetella* and *Neisseria* proliferate in Eugonic Broth because large amount of sulfur and carbon sources have been added in the formulation. Therefore Eugonic Broth is recommended for the direct isolation of *Bordetella pertussis* and *Neisseria meningitidis* from the test materials such as throat mucus, blood, cerebrospinal fluid, pleural fluid and other specimens. For the isolation of *Bacillus pumilus*, Eugonic Broth can be supplemented with 0.1% starch, prior to sterilization (3)

Tryptone and Soya peptone provide the nitrogen, vitamins and amino acids, which supports the growth of fastidious microbial species. The high concentration of dextrose is the energy source for rapid growth of bacteria. L-Cystine and sodium sulphite are added to stimulate growth. Sodium chloride maintains the osmotic balance of the media. The high carbohydrate content along with high sulfur (cystine) content improves growth with chromogenicity (4).

Type of specimen

Clinical samples - throat mucus and other specimens, Food samples

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (6,7). For food samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(5) After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

Please refer disclaimer Overleaf.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Yellow coloured, clear solution in tubes

Reaction

Reaction of 2.94% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed with added 5-10%v/v sterile defibrinated blood after an incubation at 35-37°C for 48 hours (fungal cultures incubated at 25-30°C).

Organism	Inoculum (CFU)	Growth
<i>Bacillus pumilus</i> ATCC 14884	50-100	good (with 0.1% starch)
<i>Brucella abortus</i> ATCC 4315	50-100	good (under 3-5% CO ₂)
<i>Candida albicans</i> ATCC 26790	50-100	good
<i>Lactobacillus fermentum</i> ATCC 9338	50-100	good
<i>Neisseria meningitidis</i> ATCC 13090	50-100	good
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	luxuriant (under 3-5% CO ₂)
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	luxuriant (under 3-5% CO ₂)

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

1. Pelczar and Vera J., 1949, Milk Plant Monthly 38:30
2. Niven C. F., Castellani A. G., and Allanson V., 1949, J. Bacteriol., 58:633
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams & Wilkins, Baltimore, Md.
4. Frank H. A., 1955, J. Bacteriol., 70:269.
5. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
6. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition
7. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.


Revision :04/2024



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India



**In vitro diagnostic
medical device**



Storage temperature



CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



CE Marking



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.