

# **Technical Data**

# **Disinfectant Test Broth (Staphylococcus Aureus Enrichment Broth)**

**M464** 

Disinfectant Test Broth (Staphylococcus Aureus Enrichment Broth ) is recommended for enrichment of *Staphylococcus aureus* .

# Composition\*\*

Ingredients	<b>Gms / Litre</b>
Meat (Beef) infusion	5.00
Peptic digest of animal tissue	10.000
Sodium chloride	5.000
Final pH ( at 25°C)	6.8±0.2

<sup>\*\*</sup>Formula adjusted, standardized to suit performance parameters

#### **Directions**

Suspend 20 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

# **Principle And Interpretation**

Bacteria in the genus *Staphylococcus* are pathogens of man and other mammals. Traditionally they were divided into two groups on the basis of their ability to clot blood plasma (the coagulase reaction). The coagulase-positive Staphylococci constitute the most pathogenic species Staphylococcus aureus. The presence of Staphylococci in a lesion might first be suspected after examination of a direct gram stain. However, small numbers of bacteria in blood preclude microscopic examination and require culturing first (1). Staphylococcus Aureus Enrichment Broth (Disinfectant Test Broth) is recommended for enrichment of *S.aureus* and Disinfectant testing as per ISI Specifications.

Peptic digest of animal tissue and meat infusion provide carbonaceous, nitrogenous compounds and other necessary nutrients for the growth of the test organisms. Yeast extract provides vitamins and other trace nutrients. Sodium chloride maintains the osmotic equilibrium

### **Quality Control**

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Colour and Clarity of prepared medium

Yellow coloured, clear solution without any precipitate

#### Reaction

Reaction of 2.0% w/v aqueous solution at 25°C. pH :  $6.8\pm0.2$ 

#### pН

6.60-7.00

#### **Cultural Response**

M464: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours .

Organism	Inoculum	Growth
	(CFU)	
Staphylococcus aureus	50-100	luxuriant
ATCC 25923		
Staphylococcus aureus	50-100	luxuriant
ATCC 6538		
Escherichia coli ATCC	50-100	good-luxuriant
25922		
Pseudomonas aeruginosa	50-100	good-luxuriant
ATCC 27853		

HiMedia Laboratories Technical Data

# **Storage and Shelf Life**

Store below 30°C in a tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

#### Reference

1. Easmon C. S. F., Adlam C., Staphylococci and staphylococcal infections. Vol. I & II, 1983, Academic Press, London

Revision: 2 / 2015

#### Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.