

Technical Data

BYE Agar Intended Use:

M470

Recommended for cultivation and routine studies of distribution of *Mycoplasmas* or Pleuropneumonia like organisms (PPLOs) and L-forms of bacteria.

Composition**

Ingredients	g / L
Proteose peptone	10.000
HM infusion from (200 g) #	8.000
HM infusion B from (250 g) ##	9.500
Dextrose (Glucose)	2.000
Sodium chloride	5.000
Disodium hydrogen phosphate	2.500
Yeast extract	2.000
Agar	13.000
Final pH (at 25°C)	7.9 ± 0.2

**Formula adjusted, standardized to suit performance parameters, # Equivalent to Calf brain, infusion from, ## Equivalent to Beef heart, infusion from

Directions

Suspend 52.0 grams in 850 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add 150 ml of sterile human or animal blood or serum. Mix gently and pour into sterile Petri plates.

Principle And Interpretation

Mycoplasmas (mollicutes) are the smallest free-living microorganisms (1). Earlier *Mycoplasmataceae* were given the general title of pleuropneumonia like organism (PPLO), because of similarities to *Mycoplasma mycoides* (subsp. mycoides), the causative agent of bovine pleuropneumonia (2). BYE media are simple media developed for cultivation and routine studies of distribution, habitat and possible pathogenesis of Mycoplasma - Pleuropneumonia like organisms and L-forms of bacteria by Barile, Yaguchi and Eveland (1). These media can be used for isolation of PPLOs from urethritis, penile ulcerations and cervical specimens and L-forms of *Corynebacterium, Neisseria*, and *Streptococcus*. These are also used for detecting PPLO contamination of tissue culture and cell-lines (3) and for membrane filter work (4).

BYE Agar contains Proteose peptone, HM infusion from, HM infusion B from along with yeast extract, which provide carbon, nitrogen, vitamins and other growth factors required for the metabolism of Mycoplasma - Pleuropneumonia like organisms. Inoculations are made in duplicates. One set is incubated aerobically while the other anaerobically for 48 hours or more. Usually growth occurs within 3-5 days; however, negative results are reported after 10 days. Anaerobic conditions are most important for the first 3 days while secondary transfers can be incubated aerobically.

Type of specimen

Clinical samples - Swabs of Sputum, Tissue, Bronchial lavage (BAL) fluid, Bronchial washings, Cerebrospinal fluid (CSF)

Specimen Collection and Handling:

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (5,6). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium. 2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.3% Agar gel.

Colour and Clarity of prepared medium

Yellow coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 5.2% w/v aqueous solution at 25°C. pH : 7.9±0.2

pН

7.70-8.10

Cultural Response

Cultural characteristics observed with added serum under humidified anaerobic conditions, after an incubation at 35-37°C

for 5-10 days.	
Organism	Growth
<i>Mycoplasma bovis</i> ATCC 25523	good-luxuriant
<i>Mycoplasma gallinarium</i> ATCC 19708	good-luxuriant
<i>Mycoplasma pneumoniae</i> ATCC 15531	good-luxuriant
Streptococcus pneumoniae	good-luxuriant

ATCC 6303

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

1.Murray P.R., Baron E. J., Pfaller M.A., Tenover F.C., Yolken R.H.(Eds.),1995, Manual of Clinical Microbiology, 6th Ed., ASM Press.

2.Barile, Yaguchi, Eveland, 1958, Am. J. Clin. Path. 30:171.

3.Collee J.G, Fraser A.G., Marmion B.P., Simmons. A (Eds.), 1996, Mackie and McCartney Practical Medical Microbiology, 14th Ed, Churchill Livingstone.

4.Barile, 1962, J. Bacteriol., 83:430.

5. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.

6.Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.



HiMedia Laboratories Pvt. Limited, Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area,



Thane (W) -400604, MS, India

CEpartner4U, Esdoornlaan 13, 3951DB Maarn, NL www.cepartner4u.eu



In vitro diagnostic

medical device

IVD



-30°C Storage temperature

> Do not use if package is damaged

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

HiMedia Laboratories Pvt. Ltd. Corporate Office : Plot No.C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) - 400604, India. Customer care No.: 022-6147 1919 Email: techhelp@himedialabs.com Website: www.himedialabs.com