



Buffered Charcoal Yeast Extract Agar Base

M813

Intended Use:

Recommended for selective isolation and cultivation of *Legionella* species from cooling towers, clinical and other specimens.

Composition**

Ingredients	g / L
Yeast extract	10.000
Charcoal activated	2.000
ACES buffer	10.000
α -Ketoglutarate monopotassium salt	1.000
Agar	17.000
Final pH (at 25°C)	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 20.0 grams in 500 ml purified/distilled water. Add 2.4 grams of KOH pellets and mix to dissolve. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add sterile rehydrated contents of 1 vial each of MWY Selective Supplement (FD040) and Legi Growth Supplement w/o SS (Twin Pack) (FD041A). Mix well and pour with constant stirring into sterile Petri plates, taking care that charcoal particles get evenly distributed. For additional selectivity, VCAT Selective Supplement (FD017), C3V Selective Supplement (FD037) and PGV Selective Supplement (FD038) may be added to molten medium as per choice.

Principle And Interpretation

Feeley et al (1) originally formulated Charcoal Yeast Extract (CYE) Agar. This medium was a modification of the existing F-G Agar (2). F-G Agar had starch and casein enzymic hydrolysate as ingredients in the composition. Feeley et al (1,2) replaced these two with charcoal and yeast extract respectively, and reported better recovery of *Legionella pneumophila*. Later Paseulle (3) reported that supplementation of the Charcoal Yeast Agar with ACES buffer improved the performance of the medium. Edelstein (4) further modified the medium by adding alpha-ketoglutarate. This addition helped in improving the sensitivity of the medium. Buffered Charcoal Yeast Extract Agar Base is based on Edelsteins Modification. *Legionella* species are non-spore forming, narrow, gram-negative rods. *Legionella* causes pneumonia (Legionnaires disease) (5) or a milk, febrile disease (Pontiac fever). They do not oxidize or ferment carbohydrates in conventional media or grow on sheep blood agar. Growth is much better and more rapid on Buffered Charcoal Yeast Extract Agar (2,6). Amino acids are the major sources of energy for *Legionella*. The amino acid L-cystine holds an absolute requirement as it plays major role in growth metabolism of *Legionella* (7). This amino acid as well as ferric pyrophosphate helps for the growth of *Legionella*. The media contains charcoal, which acts as a detoxicant. Yeast extract acts as a rich source of vitamins, nitrogen as well as carbon. ACES Buffer maintains optimal pH for growth while L-cystine hydrochloride; ferric pyrophosphate and α -ketoglutarate stimulate growth of *Legionella* species. For selective isolation, antibiotic supplements can be used to suppress contaminating microorganisms. C3V Selective Supplement (FD037) containing cephalothin, colistin, vancomycin and cycloheximide (8) or MWY Selective Supplement (FD040) containing glycine, polymyxin B, anisomycin, vancomycin, bromothymol blue and bromocresol purple (9) are often used. Wear gown, mask and gloves while handling *Legionella* cultures. Work in a safety hood.

Type of specimen

Clinical samples - Urine; Water samples

Specimen Collection and Handling:

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (10,11).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards (12). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic use. For professional use only. Read the label before opening the container. Wear protective gloves/ protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Further biochemical confirmation has to be carried out for further confirmation.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Grey to black homogeneous free flowing powder

Gelling

Firm, comparable with 1.7% Agar gel.

Colour and Clarity of prepared medium

Grey-black coloured opalescent gel forms in Petri plates.

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C. pH : 6.9±0.2

pH

6.70-7.10

Cultural Response

Cultural characteristics observed in 90% humid atmosphere with added MWY Selective Supplement (FD040) and Legi Growth Supplement w/o SS (Twin Pack) (FD041A), after an incubation at 35-37°C for 3-4 days.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	none-poor	≤10%	
<i>Legionella dumoffii</i> ATCC 33343	50-100	luxuriant	≥50%	light blue-grey
<i>Legionella pneumophila</i> ATCC 33153	50-100	luxuriant	≥50%	white grey to blue grey
<i>Staphylococcus epidermidis</i> ATCC 12228 (00036*)	50-100	none-poor	≤10%	

Key : (*) - Corresponding WDCM numbers

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (10,11).

Reference

1. Feeley J. C., Gibson R. J., Gorman G. W. et al, 1979, J. Clin. Microbiol., 10:437.
2. Feeley J. C., Gorman G. W., Weaver R. E. et al, 1978, J. Clin. Microbiol., 8 : 320-325.
3. Paeulle, Feely et al, 1980, J. Infect. Dis., 191:727.
4. Edelstein P. H., 1981, J. Clin. Microbiol., 14:298.

5. Broome C. V., Fraser D. W., 1979, Epidemiol. Rev 1:1-16.
6. Jones G. T., Hebert G. A., (Eds.), 1979, US Department of Health, Education and Welfare Publication No. (CDC) 8375, Atlanta, Centers for Disease Control.
7. George J. R. et al, 1980, J. Clin. Microbiol., 11:286
8. Bopp C. A., Sumner J. W., Morris G. K. and Wells J. G., 1981, J. Clin. Microbiol., 13:714.
9. Vicker R., Brown and Garrity, 1981, J. Clin. Microbiol., 13:380.
10. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
11. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W.(2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
12. Lipps WC, Braun-Howland EB, Baxter TE, eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.


Revision :03/2024



HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India



**In vitro diagnostic
medical device**



Storage temperature



CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



CE Marking



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.