



Technical Data

A C Broth

M875

Intended Use:

Recommended for cultivation of common aerobes and sterility testing of solutions and biological products without mercurial preservatives.

Composition**

Ingredients	Gms / Litre
Proteose peptone	20.000
HM peptone B #	3.000
Yeast extract	3.000
Malt extract	3.000
Dextrose (Glucose)	5.000
Ascorbic acid	0.200
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Equivalent to Beef extract

Directions

Suspend 34.2 grams in 1000 ml of purified / distilled water. Heat if necessary to dissolve the medium completely. Distribute in tubes or bottles to give the desired depth and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

If the medium is not used on same day, it is advisable to drive off dissolved gases by boiling or steaming in the autoclave and cool without agitation.

Principle And Interpretation

AC Broth support an early and luxuriant growth of aerobic, anaerobic and microaerophilic microorganisms. Many pathogenic and saprophytic aerobes can also be cultivated using AC Broth (4). This medium can also be used for sterility testing of solutions and biological products not containing mercurial preservatives. Some of the media containing sodium thioglycollate exhibit toxicity for some organisms. This toxicity is not seen in the case of AC Broth as reported by Christensen (1) and Malin and Finn (5). Earlier studies performed have reported the usefulness of using this medium for the cultivation of a wide variety of organisms (6,7).

Proteose peptone, HM peptone B, yeast extract and malt extract serve as the carbon and nitrogen sources in addition to being a source of vitamins and cofactors. Dextrose serves as the fermentable carbohydrate and source of energy. Ascorbic acid in the media helps to improve the clarity of the medium.

Type of specimen

Pharmaceutical samples- Environmental monitoring

Specimen Collection and Handling

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. This medium is general purpose medium and may not support the growth of fastidious organisms.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Medium amber coloured clear to slightly opalescent solution.

Reaction

Reaction of 3.42% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours (*Clostridium* species incubated anaerobically).

Organism	Inoculum (CFU)	Growth
<i>Clostridium perfringens</i> ATCC 12919	50-100	luxuriant
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	luxuriant
<i>Neisseria meningitidis</i> ATCC 13090	50-100	luxuriant
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	luxuriant
<i>Streptococcus mitis</i> ATCC 9811	50-100	luxuriant
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	luxuriant

Key : (*) Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

Reference

1. Christensen, 1944, Paper read at New York Meeting, American Public Health Association.
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
4. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I. Williams & Wilkins, Baltimore, Md.

5. Malin and Finn, 1951, J. Bacteriol., 62:349.
6. Reed and Orr, 1943, J. Bacteriol., 45:309.
7. Schneiter, Dunn and Caminita, 1945, Public Health Rep., 60:789.

Revision : 03/2020

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.