



Listeria Enrichment Medium Base (UVM Medium)

M890A

Intended use

Recommended for selective isolation and cultivation of *Listeria monocytogenes* from clinical specimens.

Composition**

Ingredients	g / L
Tryptone	5.000
Proteose peptone	5.000
HM peptone B #	5.000
Yeast extract	5.000
Sodium chloride	20.000
Potassium dihydrogen phosphate	1.350
Disodium hydrogen phosphate	12.000
Esculin	1.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

- Equivalent to Beef extract.

Directions

Suspend 27.17 grams in 500 ml purified/distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add rehydrated contents of 1 vial of UVM Selective Supplement I (FD136) for primary enrichment or 1 vial of UVM Selective Supplement II (FD137) for secondary enrichment. Mix well and dispense into tubes or flasks as desired.

Principle And Interpretation

Listeriosis is caused by *Listeria monocytogenes*, a short gram-positive non-sporulating rod. The bacilli are commonly found in soil and in the intestines of many animals including birds, fish, barnyard animals, dairy cattle and household pets. It is transmitted to humans by foods contaminated with faecal matter, as well as by the consumption of animal foods contaminated with the bacilli (1). Listeria Enrichment Medium Base is used for the selective cultivation and isolation of *L. monocytogenes* from clinical samples. The medium was originally formulated by Donnelly and Baigent (2). It was later modified by decreasing the nalidixic acid concentration in the selective supplements and subsequently increasing the acriflavin concentration (3). University of Vermont Modification Medium (UVM) used a two-step selective enrichment medium resulting in a higher isolation rate of *L.monocytogenes* from meat products within 3-4 days. This UVM Broth is recommended as a primary enrichment broth for recovery of heat-injured *Listeria* (4). Tryptone, proteose peptone, HM peptone B and yeast extract provide nitrogenous and carbonaceous compounds, long chain amino acids and other necessary nutrients while esculin offers differential properties to the medium. Nalidixic acid and acriflavin hydrochloride together with higher concentration of phosphate render the medium selective for *Listeria*. Gram-negative and gram-positive organisms are inhibited by nalidixic acid and acriflavin hydrochloride respectively. The two-step selective enrichment method developed (2) results in a higher detection rate of *L.monocytogenes* from specimens and has the added advantage of only taking 3-4 days. For primary isolation inoculate 25 gm or 25 ml specimen in 225 ml Listeria Enrichment Medium Base with added UVM Selective Supplement I (FD136). After 24 hours incubation, spread 0.2 ml of this medium on Listeria Selective Agar (M567) plate. Simultaneously transfer 0.1 ml of Enrichment broth to 10 ml of fresh Listeria Enrichment Medium Base with added UVM Selective Supplement II (FD137). For secondary enrichment after 24 hours spread 0.2 ml of this medium on Listeria Selective Agar (M567) plate. Note: Broth cultures of *Listeria* are more dangerous than colonies on agar plates, so proper precautions should be taken while handling.

Type of specimen

Clinical samples

Specimen Collection and Handling:

Please refer disclaimer Overleaf.

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (5,6).
After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. For professional use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets

Limitations :

1. The medium is not differential, so further biochemical testing is required for identification between *Listeria* species.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to light tan homogeneous free flowing powder

Colour and Clarity of prepared medium

Medium amber coloured, slightly opalescent solution with a bluish tinge

Reaction

Reaction of 5.43% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours .

Organism	Inoculum (CFU)	Growth(On addition of FD136 or FD137)
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	none to poor
<i>Listeria monocytogenes</i> ATCC 19111 (00020*)	50-100	good-luxuriant
<i>Listeria monocytogenes</i> ATCC 19112	50-100	good-luxuriant
<i>Listeria monocytogenes</i> ATCC 19117	50-100	good-luxuriant
<i>Listeria monocytogenes</i> ATCC 19118	50-100	good-luxuriant
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	none to poor

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

1. Alcamo E. I., 2001, Fundamentals of Microbiology, 6th Edition, Jones and Bartlett publishers.
2. Donnelly C. W. and Baigent G. J., 1986, Appl. Environ. Microbiol., 52:689.
3. McClain D. and Lee W. H., 1988, J. Assoc. off Anal. Chem., 71:660.
4. Bailey J. S., Fletcher D. L. and Cox N. A., 1990, J. Food Prot., 53:473.
5. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.


Revision : 03/2024




HiMedia Laboratories Pvt. Limited,
Plot No.C-40, Road No.21Y,
MIDC, Wagle Industrial Area,
Thane (W) -400604, MS, India



**In vitro diagnostic
medical device**



Storage temperature



CEpartner4U, Esdoornlaan 13,
3951DB Maarn, NL
www.cepartner4u.eu



CE Marking



**Do not use if
package is damaged**

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.