



Technical Data

Luria Bertani HiCynth™ Agar, Miller (Miller Luria Bertani HiCynth™ Agar) MCD1151

Luria Bertani HiCynth™ Agar, Miller is used for the cultivation and maintenance of recombinant strains of *Escherichia coli* for genetic and molecular studies and may be used for routine cultivation of not particularly fastidious microorganisms.

Composition**

Ingredients	Gms / Litre
HiCynth™ Peptone No.2*	10.000
HiCynth™ Peptone No.5*	5.000
Sodium chloride	10.000
Agar	15.000
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

*Chemically defined peptones

Directions

Suspend 40 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Luria Bertani Agar is prepared as described by Lennox (1) for cultivation and maintenance of recombinant strains of *Escherichia coli*. Luria Bertani Agar, Miller (2) is slightly different with double amount of sodium chloride. Luria Bertani HiCynth™ Agar, Miller is a modification of Luria Bertani Agar, Miller wherein animal based peptones are replaced by chemically defined peptones to avoid BSE/TSE risks associated with animal based peptones. The media is nutritionally rich for the growth of pure cultures of recombinant strains. Strains derived from *Escherichia coli* K12 are deficient in Vitamin B synthesis, are further modified by specific mutation to create auxotrophic strains and are therefore unable to grow on nutritionally deficient media.

HiCynth™ Peptone No.2 and HiCynth™ Peptone No.5 provides nitrogenous and carbonaceous compounds, amino acids, vitamins and other growth factors required for the growth. Sodium chloride provides sodium ions for membrane transport and also maintains the osmotic equilibrium of the medium.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow to amber coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C. pH : 7.5±0.2

pH

7.30-7.70

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18 - 24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response <i>Escherichia coli</i> ATCC 23724	50-100	luxuriant	>=70%

Please refer disclaimer Overleaf.

<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=70%
<i>Escherichia coli</i> DH5 alpha MTCC 1652	50-100	luxuriant	>=70%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label.

Reference

- 1.Lennox E.S., 1955, Transduction of Linked Genetic Characters of the host by bacteriophage P1., Virology, 1:190.
- 2.Atlas R.M., 1993, Handbook of Microbiological Media, Ed. by Parks L., CRC Press, Inc.

Revision : 00 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.