

Violet Red Bile Glucose Agar

MH581

Intended use

Recommended for detection and enumeration of *Enterobacteriaceae* from pharmaceutical products in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP .

Composition**

| Ingredients | Gms / Litre |
|-----------------------------|-------------|
| Yeast extract | 3.000 |
| Gelatin peptone # | 7.000 |
| Bile salts | 1.500 |
| Sodium chloride | 5.000 |
| Glucose monohydrate | 10.000 |
| Agar | 15.000 |
| Neutral red | 0.030 |
| Crystal violet | 0.002 |
| pH after heating (at 25°C) | 7.4±0.2 |

**Formula adjusted, standardized to suit performance parameters

Pancreatic digest of gelatin

Directions

Suspend 40.62 grams (the equivalent weight of dehydrated medium per litre) in 1000 ml purified /distilled water. Heat to boiling to dissolve the medium completely. **DONOT HEAT IN AN AUTOCLAVE**. Cool to 45 - 50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Violet Red Bile Glucose Agar is a selective medium recommended for detection and enumeration of *Enterobacteriaceae* especially the bile tolerant gram negative bacteria in accordance with the microbial limit testing by harmonized methodology of USP/EP/BP/JP/IP (1,2,3,4,5) from non-sterile products and pharmaceutical preparations.

Gelatin peptone and yeast extract provide nitrogenous, carbonaceous compounds, long chain amino acids, vitamins and other nutrients essential for bacterial metabolism. This media is selective due to presence of the inhibitors; bile salts positive organisms especially Staphylococci. Neutral red indicator helps to detect glucose fermentation. Glucose fermenting and crystal violet. Crystal violet inhibits gram-strains produce red colonies with pink-red halos in the presence of neutral red. Sodium chloride maintains the osmotic equilibrium in the medium. The red colour is due to absorption of neutral red and a subsequent colour change of the dye when the pH of medium falls below 6.8.

Type of specimen

Pharmaceutical samples.

Specimen Collection and Handling

For pharmaceutical samples, follow appropriate techniques for sample collection, processing as per guidelines (1,2,3,4,5). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions:

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Though the medium is for selective isolation of *Enterobacteriaceae*, further biochemical and serological testing must be carried out for further confirmation.
2. Over incubation may result in reverting of reaction.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to pinkish beige homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Reddish purple coloured clear to slightly opalescent gel forms in Petri plates.

pH

7.20-7.60

Growth Promotion Test

Growth Promotion is carried out in accordance with the harmonized method of USP/EP/BP/JP. Cultural response was observed after an incubation at 30-35°C for 18-24 hours. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

Growth promoting properties

Growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≤ 100 cfu (at 30-35°C for ≤ 18 hours).

Indicative properties

Colonies are comparable in appearance and indication reaction to those previously obtained with previously tested and approved lot of medium occurs for the specified temperature for a period of time within the range specified inoculating ≤ 100 cfu (at 30-35°C for 18-24 hours).

Cultural Response

| Organism | Inoculum (CFU) | Growth | Observed Lot value (CFU) | Recovery | Colour of colony | Incubation temperature |
|---|----------------|----------------|--------------------------|-------------|--------------------------------|------------------------|
| Growth Promoting + Indicative | | | | | | |
| <i>Escherichia coli</i> ATCC 8739 (00012*) | 50 -100 | luxuriant | 25 -100 | ≥ 50 % | pink-red with bile precipitate | 18 -24 hrs |
| <i>Pseudomonas aeruginosa</i> ATCC 9027 (00026*) | 50 -100 | luxuriant | 25 -100 | ≥ 50 % | pink to red | 18 -24 hrs |
| Additional Microbiological Testing | | | | | | |
| <i>Escherichia coli</i> NCTC 9002 | 50 -100 | good-luxuriant | 25 -100 | ≥ 50 % | pink-red with bile precipitate | 18 -24 hrs |
| <i>Escherichia coli</i> ATCC 25922 (00013*) | 50 -100 | good-luxuriant | 25 -100 | ≥ 50 % | pink-red with bile precipitate | 18 -24 hrs |
| <i>Salmonella Enteritidis</i> ATCC 13076 (00030*) | 50 -100 | good-luxuriant | 25 -100 | ≥ 50 % | light pink | 18 -24 hrs |
| # <i>Klebsiella aerogenes</i> ATCC 13048 (00175*) | 50 -100 | good-luxuriant | 25 -100 | ≥ 50 % | pink-red | 18 -24 hrs |
| <i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*) | $\geq 10^3$ | inhibited | 0 | 0% | | ≥ 24 hrs |
| <i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*) | $\geq 10^3$ | inhibited | 0 | 0% | | ≥ 24 hrs |

Key :-(#) Formerly known as *Enterobacter aerogenes* , (*) Corresponding WDCM numbers

Please refer disclaimer Overleaf.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (6,7).

Reference

1. The United States Pharmacopoeia-National Formulary (USP-NF), 2022.
2. The British Pharmacopoeia, 2022, Medicines and Healthcare products Regulatory Agency.
3. European Pharmacopoeia, 2022, 10 th volume, European Directorate for the quality of medicines & Healthcare.
4. The Japanese Pharmacopoeia, 17th edition, 2016, The Ministry of Health, Labour and welfare.
5. Indian Pharmacopoeia, 2022, Indian Pharmacopoeia Commission, Ministry of Health and Family Welfare Government of India.
6. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
7. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W.(2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 05/2023

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.