

Modified Plate Count Agar Plate

MP163

Intended use

Recommended for enumeration of bacteria in milk and milk products, rinse waters, ice creams etc.

Composition**

Ingredients	Gms / Litre
Peptone	5.000
Yeast extract	3.000
M-protein solids#	1.000
Agar	15.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

#- Equivalent to Milk solids

Directions

Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.

Principle And Interpretation

The milk secreted in an uninfected cow's udder is sterile. Contamination of this milk can occur during milking, cooling and storage (1). Milk is an excellent medium for bacteria, yeast and moulds. Their rapid growth can cause marked deterioration, spoiling the milk for liquid consumption or manufacture into dairy products. Human infection can occur by consumption of such contaminated milk or milk products. Milk Agar is recommended for performing plate count tests on milk, rinse waters and dairy products. It is formulated as per the official medium described by Department of Health Memo (2). It is also recommended by EUROGLACE (EEC Ice Cream Industries) for the examination of ice cream (2).

Peptone and yeast extract provide essential nutrients while M-Protein solids are a source of casein. Dextrose is the carbon and energy source. Proteolytic bacteria will be surrounded by a clear zone, due to the conversion of casein into soluble nitrogenous compounds (3).

For milk, dilutions of 1/10, 1/100 and 1/1000 are prepared with 1/4 strength Ringer solution. 1 ml of each dilution is pipetted aseptically into sterile Petri plates to which 10 ml of sterile and cooled Milk Agar is added and mixed well. Plates should be poured within 15 minutes of dilution preparation. After solidification of medium the plates are allowed to stand for 1 hour before transferring to the incubator. Incubate at 35°C for 2 or 3 days at 30°C. Higher counts may be obtained after an incubation at 22°C and 30°C temperature rather than at 35°C (3,4,5). Count the colonies within 4 hours after the incubation and read it as per ml of sample.

Type of specimen

Milk and milk products, rinse waters, ice creams etc.

Specimen Collection and Handling:

For milk and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (6,7). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's requirement.
3. Further serological and biochemical testing is required for complete identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Sterile Modified Plate Count Agar in 90 mm disposable plates with smooth surface and absence of black particles/cracks/bubbles

Colour of medium

Light yellow coloured medium

pH

7.00-7.40

Cultural Response

Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Bacillus subtilis</i> subsp. <i>spizizenni</i> ATCC 6633 (00003*)	50-100	good-luxuriant	>=70%
<i>Pseudomonas aeruginosa</i> ATCC 27853 (00025*)	50-100	good-luxuriant	>=70%
<i>Lactobacillus casei</i> ATCC 9595	50-100	good-luxuriant	>=70%
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	good-luxuriant	>=70%
<i>Serratia marcescens</i> ATCC 8100	50-100	good-luxuriant	>=70%

*- Corresponding WDCM numbers

Storage and Shelf Life

On receipt store between 20-30°C. Use before expiry date on the label. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (8,9).

Reference

1. Collee J. G., Fraser A. G., Marimon B. P., Simmons A., (Eds.), Mackie and McCartney, Practical Medical Microbiology, 1996, 14th Edition, Churchill Livingstone.
2. Dept. of Health, 1987, Memo. 139/Foods.
3. Methods of Microbiological Examination for Dairy Purposes, Diluents, Media and Apparatus and their Preparation and Sterilization, BS4285, Sec. 1.2.
4. Davis J. G., 1959, Milk Testing, 2nd Ed., Dairy Industries Ltd., London, Pg. 175.
5. Klose J., 1968, Susswaren, 14:778.
6. Thomas S. B. and Jenkins E., 1940, Proc. Soc. Appl. Agric., 38:40.
7. Wilson G. S., 1935, Bacteriological Grading of Milk, HMSO, London.
8. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
9. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision :02/2023

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.