

Field's Tryptic digest Broth, HiVeg™ (Tryptic Digest Broth, HiVeg™)

MV1028

Intended Use:

Recommended for cultivation of fastidious microorganisms from various samples.

Composition**

Ingredients	g / L
HiVeg™ infusion	10.000
Sodium chloride	5.000
Dextrose (Glucose)	1.000
Final pH (at 25°C)	7.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 16.0 grams in 1000 ml purified/distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add sterile enrichment such as blood, serum or ascitic fluid as required. Mix well and dispense into sterile test tubes or flasks as required.

Principle And Interpretation

Infusion based media are routinely employed for the cultivation of fastidious organisms. These media suffice complex nutritional requirements of most of the organisms. Field (1) formulated Tryptic Digest Broth for cultivation of fastidious microorganisms. Supplementation of the medium with serum, blood or ascitic fluid, it supports the growth of fastidious organisms such as, *Neisseria meningitides*, *Haemophilus influenzae*, *Streptococcus pneumoniae* etc. (2). Shepard et al (3) described its use for culturing *Streptococcus pneumoniae* and *Actinomycetes*. Field's Tryptic digest Broth, HiVeg™ is prepared by using vegetable peptones in place of animal based peptones which make the media free of BSE/TSE risks. HiVeg™ infusion provides carbonaceous and nitrogenous compounds and other essential growth nutrients for growth of fastidious microorganisms. Blood serum or ascitic fluid provides additional growth factors.

Type of specimen

Please add specimens

Specimen Collection and Handling:

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Further biochemical and serological tests must be carried out for further identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to light yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Basal Medium : Light yellow coloured clear solution. After addition of blood, cherry red coloured and with added serum or ascitic fluid, dark yellow coloured, opaque or clear to slightly opalescent solution respectively.

Reaction

Reaction of 1.6% w/v aqueous solution at 25°C. pH : 7.6±0.2

pH

7.40-7.80

Cultural Response

Cultural characteristics observed with added blood, serum or ascetic fluid after incubation at 35-37°C for 24 - 48 hours.

Organism	Inoculum (CFU)	Growth
<i>Bordetella pertussis</i> ATCC 8467	50-100	luxuriant
<i>Haemophilus influenzae</i> ATCC 35056	50-100	luxuriant
<i>Neisseria meningitidis</i> ATCC 13090	50-100	luxuriant
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	luxuriant
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	luxuriant
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	luxuriant

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference

1. Field V. J., 1956, SAB Newsletter, 22:8.
2. Atlas R. M., 2004, Handbook of Microbiological Media, Lawrence C. Parks (Ed.), 3rd Edition, CRC Press.
3. Shepard M. C. and Lunceford C. D., 1976, J. Clin. Microbiol., 3:613.1
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision :03/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.