



Technical Data

Starch HiVeg® Agar

MV107

Intended use

Recommended for the detection of starch hydrolyzing microorganisms.

Composition**

Ingredients	g / L
HiVeg® peptone	5.000
Sodium chloride	5.000
Yeast extract	1.500
HiVeg® extract	1.500
Starch, soluble	2.000
Agar	15.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 30.0 grams in 1000 ml purified/ distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Starch Agar was formulated by Vedder (1) in 1915, for the cultivation of *Neisseria*. Since then, other media have been developed that are superior to Starch Agar for the isolation of *Neisseria* species, including enriched GC Medium Base. Starch Agar (M107) is recommended for the detection of starch hydrolyzing microorganisms from foods (2) and clinical samples (3). Although the medium was originally formulated to perform the test for the identification of *Bacillus cereus*, it can be applied to any kind of microorganism where starch hydrolysis activity is required to be analyzed. Starch HiVeg® Agar is same as Starch Agar except that the animal based peptones are completely replaced with vegetable peptones to avoid the BSE/TSE risks associated with animal peptones.

HiVeg® peptone, yeast extract and HiVeg® extract provide nitrogenous compounds, carbon, sulphur, trace elements etc. to the microorganisms. Sodium chloride maintains osmotic equilibrium. Flood the surface of 48 hours old culture on Starch HiVeg® Agar with Grams Iodine (S013). Starch hydrolysis is seen as a colourless zone surrounding the colonies. A blue or purple zone indicates that starch is not hydrolyzed. Size of the clear zone is directly proportional to the starch hydrolyzing activity of the strain under study.

Type of specimen

Isolated Microorganism

Specimen Collection and Handling:

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (4,5).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.

(6) After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheet

Limitations:

1. Due to nutritional variations, some strains may show poor growth.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Please refer disclaimer Overleaf.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of Prepared Medium

Yellow coloured slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.0% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours. (\$) - on addition of Iodine solution)

Organism	Inoculum (CFU)	Growth	Starch hydrolysis \$
** <i>Bacillus spizizenii</i> ATCC 6633(00003*)	50-100	luxuriant	positive reaction ,clearing around the colony
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	luxuriant	negative reaction
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00032*)	50-100	luxuriant	negative reaction
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	luxuriant	negative reaction

*- Corresponding WDCM numbers

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

Reference

1. Vedder E. B., 1915, J. Infect. Dis., 16:385.
2. Harrigan W. and McCance M., 1976, Laboratory Methods in Food and Dairy Microbiology, Academic Press Inc. (London) Ltd.
3. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Eds.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
4. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
5. Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
6. Lipps WC, Braun-Howland EB, Baxter TE, eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.

7. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
8. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.385.

Revision : 02 / 2025

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.