



# Technical Data

## Rapid HiColiform™ HiVeg™ Agar

MV1465

### Intended use

Recommended for detection and confirmation of *Escherichia coli* and total coliforms on the basis of enzyme substrate reaction from water samples, using a combination of chromogenic and fluorogenic substrate.

### Composition\*\*

Ingredients	g / L
HiVeg™ special peptone	5.000
Sodium chloride	5.000
Sorbitol	1.000
Dipotassium hydrogen phosphate	2.700
Potassium dihydrogen phosphate	2.000
Sodium lauryl sulphate	0.100
Chromogenic substrate	0.080
Fluorogenic substrate	0.050
IPTG(1-Isopropyl-β-D-1-thiogalactopyranoside)	0.100
Agar	15.000
Final pH ( at 25°C)	6.8±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 31.03 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

### Principle And Interpretation

The Rapid HiColiform™ Agar is modification of LMX Broth described by Manafi and Kneifel (1). Rapid HiColiform™ Agar is used for the simultaneous detection of total coliforms and *Escherichia coli*. This media is useful for the detection and confirmation of *Escherichia coli* and total coliforms in water samples on the basis of chromogenic and fluorogenic substrates (2,1,3,4,5,6). It can also be used for clinical samples.

Rapid HiColiform™ HiVeg™ Agar is prepared by completely replacing animal peptones with vegetable peptone to avoid BSE/TSE risks. HiVeg™ special peptone which is rich in tryptophan content, provides essential growth nutrients and is useful for the simultaneous detection of indole production. Sorbitol provides the carbon source. The phosphate salts provide buffering action for rapid growth of coliforms. Sodium lauryl sulphate makes the medium selective by inhibiting accompanying microflora, especially the gram-positive organisms. The fluorogenic substrate, is split by enzyme β-D-glucuronidase, which is specifically found in *Escherichia coli*. The reaction is indicated by a blue fluorescence under UV light. The presence of total coliforms is indicated by a blue-green colour of the colonies due to the cleavages of the chromogenic substrate. IPTG amplifies enzyme synthesis and increases the activity of β-D-galactosidase. To confirm presence of *E. coli*, add 2-3 drops of Kovacs reagent over the suspected colony. The colony turns red within 2 minutes in case of positive reaction.

### Type of specimen

Food samples; Water samples.

### Specimen Collection and Handling:

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards (7). After use, contaminated materials must be sterilized by autoclaving before discarding.

### Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

## Limitations :

1. 97% of the *E.coli* strains are  $\beta$ -D-glucuronidase positive, however few strains may show negative fluorescence.

## Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Cream to yellow homogeneous free flowing powder

### Gelling

Firm, comparable with 1.5% Agar gel

### Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates

### Reaction

Reaction of 3.1% w/v aqueous solution at 25°C. pH : 6.8±0.2

### pH

6.60-7.00

### Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Colour of medium/ Colony	Fluorescence (under uv)	Indole production
# <i>Klebsiella aerogenes</i> ATCC 13048 (00175*)	50-100	luxuriant	blue-green	negative reaction	negative reaction
<i>Escherichia coli</i> ATCC 25922 (00012*)	50-100	luxuriant	blue-green	positive reaction	positive reaction
<i>Klebsiella pneumoniae</i> ATCC 13883 (00097*)	50-100	luxuriant	blue-green	negative reaction	negative reaction
<i>Salmonella</i> Typhimurium ATCC 14028 (00031*)	50-100	luxuriant	yellow	negative reaction	negative reaction

# Formerly known as *Enterobacter aerogenes*

Key : (\*) Corresponding WDCM numbers.

## Storage and Shelf Life

Store between 15-25°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (8,9).

## Reference

1. Manafi. M. and Kneifel W., (1989), Zbl. Hygiene and Umweltmedizin 189:225-234.
2. Hahn G. and Wittrock E., (1991), Acta Microbiologica Hungarica 38(3-4):265-271.
3. Manafi M., (1990), Forum Stadte-Hygiene 41:181-184.
4. Manafi M., (1991), Ernährung / Nutrition, 15, Nr. 10.
5. Manafi M. and Kneifel W., (1991), Acta Microbiologica Hungarica 33(3-4):293-304.
6. Manafi M., Kneifel B. and Bascon S., (1991), Microbiol. Rev., 55:335-348.
7. Lipps WC, Braun-Howland EB, Baxter TE, eds. Standard methods for the Examination of Water and Wastewater, 24th ed. Washington DC:APHA Press; 2023.

8. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
9. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 03 / 2024

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.