

## BHI CC Agar, HiVeg™

MV209

### Intended Use:

Recommended for selective isolation and cultivation of fastidious pathogenic fungi from specimens heavily contaminated with bacteria and saprophytic fungi.

### Composition\*\*

Ingredients	g / L
HiVeg™ special infusion	7.50
HiVeg™ infusion	10.000
HiVeg™ peptone No. 3	10.000
Dextrose (Glucose)	2.000
Sodium chloride	5.000
Disodium hydrogen phosphate	2.500
Chloramphenicol	0.050
Cycloheximide	0.500
Agar	15.000
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 52.5 grams in 1000 ml purified / distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Avoid excess heat as it may reduce the selectivity of the medium. Mix well and pour into sterile Petri plates.

Warning : Cycloheximide is very toxic. Avoid skin contact or aerosol formation and inhalation.

### Principle And Interpretation

Brain Heart CC Agar is formulated as per Ajello et al. (1) and McDonough, et al. (2). This medium is recommended for selective isolation of pathogenic fungi (3). Chloramphenicol is a broad-spectrum antibiotic, which inhibits the growth of wide range of gram-positive and gram-negative bacteria. Cycloheximide inhibits most saprophytic moulds and enhances the isolation of pathogenic fungi.

BHI CC Agar, HiVeg™ is prepared by using vegetable peptones in place of animal based peptones which make the media free of BSE/TSE risks.

This medium contains HiVeg™ special infusion and HiVeg™ infusion infusion and HiVeg™ peptone No. 3 to supply the necessary nutrients to support the growth of fastidious pathogenic fungi. Dextrose is a carbohydrate source and disodium phosphate buffers the medium. The medium may be further enriched with 10% sheep blood to isolate systemic fungi that grow poorly on non-enriched medium. Also the addition of Gentamicin, 50 mcg/ml of medium, improves the selectivity. The antibiotics in this medium may inhibit some fungi. The addition of blood makes BHI CC Agar suitable for the isolation of the tissue phase of *Histoplasma capsulatum* and other pathogenic fungi, including *Coccidioides immitis*.

While handling *Histoplasma capsulatum* extreme care should be taken to avoid dissemination of its infective spores. The culture should be examined in a closed filtered air cabinet. Isolation of fungi from contaminated specimens can be done by inoculating selective medium along with nonselective medium and incubated at 25-30°C. For isolation of fungi causing systemic mycoses two sets of media should be inoculated with one set incubated at 23-30°C and a duplicate set at 35-37°C. Examine the plates for at least a week.

### Type of specimen

Please add specimens

### Specimen Collection and Handling

After use, contaminated materials must be sterilized by autoclaving before discarding.

## Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

## Limitations

1. Further biochemical tests must be carried out for complete identification.

## Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Cream to yellow homogeneous free flowing powder

### Gelling

Firm, comparable with 1.5% Agar gel

### Colour and Clarity of prepared medium

Light amber coloured, clear to slightly opalescent gel forms in Petri plates.

### Reaction

Reaction of 5.25% w/v aqueous solution at 25°C. pH : 7.4±0.2

### pH

7.20-7.60

### Cultural Response

Cultural characteristics observed after an incubation at 25-35°C for 40-96 hours. (*Trichophyton* species incubated for 102 weeks)

Organism	Growth
<i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	inhibited
<i>Blastomyces dermatidis</i> ATCC 14112	good
<i>Candida tropicalis</i> ATCC 1369	inhibited
<i>Candida albicans</i> ATCC 26790	fair-good
<i>Escherichia coli</i> ATCC 25922 (00013*)	inhibited
<i>Histoplasma capsulatum</i> ATCC 10230	good
<i>Trichophyton megninii</i> ATCC 12106	good-luxuriant
<i>Trichophyton mentagrophytes</i> ATCC 9533	good-luxuriant
<i>Trichophyton tonsurans</i> ATCC 10220	good-luxuriant

Key : \*Corresponding WDCM numbers.

## Storage and Shelf Life

Store dehydrated and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

---

## Reference

1. Ajello L., George L., Kaplan W., and Kaufman L., 1966, CDC Laboratory Manual of Medical Mycology, Atlanta, Ga: US. DHEW, Center for Disease Control.
2. McDonough E., George L., Ajello L., and Brinkman S., 1960, Mycopathol. Mycol. Appl; 13:113.
3. Murray P. R., Baron E. J., Jorgensen J. H., Pfaller M. A., Tenover F. C., and Tenover F. C. (Eds.), 8th ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 03 / 2024

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.