

Marine Oxidation Fermentation HiVeg™ Medium (MOF HiVeg™ MV379 Medium)

Intended Use:

Recommended for differentiation of marine bacteria on the basis of fermentative and oxidative metabolism of carbohydrates.

Composition**

Ingredients	g / L
HiVeg™ hydrolysate	1.000
Yeast extract	0.100
Tris (hydroxymethyl) aminomethane	0.500
Boric acid	0.011
Ammonium sulphate	0.500
Disodium hydrogen phosphate	0.004
Ammonium nitrate	0.0008
Sodium chloride	9.700
Magnesium chloride	4.400
Sodium sulphate	1.600
Calcium chloride anhydrous	0.900
Potassium chloride	0.275
Sodium bicarbonate	0.080
Potassium bromide	0.040
Strontium chloride	0.017
Sodium silicate	0.002
Sodium fluoride	0.0012
Phenol red	0.010
Agar	3.000
Final pH (at 25°C)	8.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 22.14 grams in 1000 ml purified / distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add sterile dextrose solution (or other carbohydrate of choice) to a final concentration of 1%. Mix well and dispense into sterile test tubes to form butts.

Principle And Interpretation

Some organisms metabolize glucose oxidatively and others ferment glucose fermentatively when the hydrogen acceptor is not oxygen. Such organisms can be differentiated based on the Oxidation Fermentation Test. This test is also known as the “oxferm” test. MOF medium is a modified version of the formula originally developed by Leifson (1); used for differentiating oxidative and fermentative carbohydrate metabolizing marine bacteria. The marine environment of the oceans illustrates a different view of microbial populations in water. In the high salt concentration of ocean water, halophilic or salt-loving microorganisms survive. In addition, the organisms must be psychrophilic since it is very cold below the surface. Those at bottom must also withstand great pressure and are therefore barophilic or pressure loving (2). HiVeg™ hydrolysate and yeast extract in the medium supply the necessary nitrogenous nutrients including amino acids, vitamins etc. The mineral content of this medium is equivalent to one-half that of seawater (1). It contains a variety of salts found in seawater, which not only makes the medium suitable for marine bacteria but also buffers the medium. Phenol red is the pH indicator in the medium. MOF HiVeg™ Medium is prepared by using vegetable peptones in place of animal based peptones which make the media free of BSE/TSE risks. For differentiating the fermentation and oxidation of carbohydrates, inoculate two tubes of medium containing carbohydrate with each culture to be tested. Cover one medium tube of each culture with sterile melted petrolatum to form a layer of about one inch in height.

Carbohydrate -fermenting marine bacteria change the colour of the medium in both the tubes (covered and uncovered) from red to yellow whereas carbohydrate-oxidizing marine bacteria change the colour of the medium from red to yellow only in the uncovered (open) tube. Marine bacteria that are neither oxidative nor fermentation do not exhibit any change in the covered medium and exhibit an alkaline (red to deep pink) reaction in the uncovered medium. Gas production is detected as splitting or displacement of agar or formation of small bubbles. Motile organisms form a diffuse zone of growth originating from the line of inoculation. Non-motile organisms grow along the line of inoculation.

Type of specimen

Isolated Microorganism from Water sample.

Specimen Collection and Handling:

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(3)
After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

N.A.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to pinkish purple homogeneous free flowing powder

Gelling

Semisolid, comparable with 0.3% Agar gel.

Colour and Clarity of prepared medium

Pink coloured clear to slightly opalescent gel forms in tubes as butts

Reaction

Reaction of 2.2% w/v aqueous solution at 25°C. pH : 8.0±0.2

pH

7.80-8.20

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Motility	Acid	Gas
<i>Vibrio cholerae</i> ATCC 15748	50-100	luxuriant	positive, growth away from stabline causing turbidity	positive reaction, yellow colour	positive reaction
<i>Vibrio parahaemolyticus</i> ATCC 17802 (00037*)	50-100	luxuriant	negative, growth along the stabline, surrounding medium remains clear	negative reaction	negative reaction

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Please refer disclaimer Overleaf.

Reference

1. Leifson E., 1963, J. Bacteriol., 85:1183.
2. Alcamo E. I., 2001, Fundamentals of Microbiology, 6th Ed., Jones and Bartlett Publishers.
3. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 03/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.