

RS HiVeg™ Medium Base

MV576

Intended use

Recommended for selective isolation, cultivation and presumptive identification of *Aeromonas hydrophila*.

Composition**

Ingredients	Gms / Litre
Yeast extract	3.000
Maltose	3.500
L-Cysteine hydrochloride	0.300
L-Lysine hydrochloride	5.000
L-Ornithine hydrochloride	6.500
Sodium thiosulphate	6.800
Ferric ammonium citrate	0.800
Synthetic detergent No. III	1.000
Sodium chloride	5.000
Bromothymol blue	0.030
Agar	13.500
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 45.43 grams in 990 ml purified/distilled water. Heat to boiling to dissolve the medium completely. **DO NOT AUTOCLAVE**. Cool to 45-50°C and aseptically add rehydrated content of 1 vial of NO 5 Selective Supplement (FD096). Mix well and pour into sterile Petri plates.

Principle And Interpretation

RS Medium was formulated by Rimler and Shotts (1) based on the principle of Xylose-Lysine (XL) Agars (2,3). It is used for selective isolation and presumptive identification of *Aeromonas hydrophila* and other gram-negative bacteria based on their ability to decarboxylate lysine and ornithine, maltose fermentation and H₂S production (4). Rimler-Shotts (RS) HiVeg™ Medium Base is prepared by completely replacing animal based peptones with vegetable peptones to avoid BSE/TSE risks associated with animal peptones.

Yeast extract acts as a source of nutrients. Sodium thiosulphate, L-cysteine hydrochloride and ferric ammonium citrate are the indicators of H₂S production. The medium contains maltose, which is mostly fermented by all *Aeromonas*. Maltose fermentation is indicated by bromothymol blue. Synthetic detergent No. III and novobiocin inhibit gram-positive bacteria and *Vibrio* species. *Citrobacter freundii* usually produce H₂S but occasionally negative strains exist. The medium contains L-cysteine and L-ornithine, which are often decarboxylated by enteric bacteria to give alkaline products. Lysine positive and ornithine positive strains of *Aeromonas* may not have the typical strong yellow colour because of alkaline products produced during decarboxylation of the amino acids. Results are interpreted within 24 hours since after 26 hours slow reversion of yellow colour to a basic (green) colour occurs. Medium should be incubated at 35°C, which will eliminate possible growth of *Aeromonas salmonicida*, which may grow at reduced temperatures giving false-positive reaction. Test the yellow colonies with or without black centers (of H₂S) for oxidase to rule out *Citrobacter* species. *Proteus mirabilis* is inhibited on this medium.

Type of specimen

Food samples ; Water samples.

Specimen Collection and Handling:

For food samples, follow appropriate techniques for sample collection and processing as per guidelines (5).

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards (6).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens.

Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Pure isolate must be used.
2. Results are interpreted within 24 hours since after 26 hours slow reversion of yellow colour to a basic (green) colour occurs.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Light yellow to light green homogeneous free flowing powder

Gelling

Firm, comparable with 1.35% Agar gel.

Colour and Clarity of prepared medium

Dark green coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.54% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH

6.80-7.20

Cultural Response

Cultural characteristics observed with added NO 5 Selective Supplement(FD096) after an incubation at 35-37°C for 24 hours

Organism	Inoculum (CFU)	Growth	Maltose fermentation	Lysine/ Ornithine decarboxylation	H ₂ S
<i>Aeromonas hydrophila</i> ATCC 7966 (00063*)	50-100	good	positive reaction, yellow coloured colonies	negative reaction	negative reaction
<i>Citrobacter freundii</i> ATCC 8090	50-100	good	negative reaction	variable reaction	positive, black centered colonies
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	good	negative reaction	variable reaction	negative reaction
## <i>Proteus hauseri</i> ATCC 13315	50-100	good	positive reaction, yellow coloured colonies	negative reaction	positive, black centered colonies
<i>Salmonella</i> Typhi ATCC 6539	50-100	good	positive reaction, yellow coloured colonies	negative reaction	negative reaction

Key : (*) Corresponding WDCM numbers ## - Formerly known as *Proteus vulgaris*

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

References

1. Shotts E. B. Jr. and Rimler R., 1973, Appl. Microbiol., 26(4):550.
2. Taylor W. I. and Harris B., 1965, Am. J. Clin. Pathol., 44:476.
3. Taylor W. I., 1965, Am. J. Clin. Pathol., 44:471.
4. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
5. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
6. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
7. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
8. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 05/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.