

## L.D. Esculin HiVeg™ Agar

MV743

### Intended Use:

Recommended for identification of anaerobic bacteria especially *Bacteroides* species on the basis of Esculin hydrolysis.

### Composition\*\*

Ingredients	g / L
HiVeg™ hydrolysate	5.000
Yeast extract	5.000
Sodium chloride	2.500
L-Tryptophan	0.200
Vitamin K1	0.010
L-Cystine	0.400
Ferric pyrophosphate	0.010
Esculin	1.000
Ferric citrate	0.500
Agar	20.000
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 34.62 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

### Principle And Interpretation

Organisms that grow in the absence of oxygen are termed as anaerobes. Depending upon their ability to tolerate oxygen, they are classified as either facultative or obligate anaerobes. The anaerobic gram-negative bacteria are part of the normal flora of the upper respiratory tract, mouth, intestinal tract and urinogenital tract of human and animals. The bile-resistant *Bacteroides fragilis* group is the most commonly recovered anaerobe in clinical specimens and is more resistant to antimicrobial agents than any other anaerobe. *Fusobacterium necrophorum* is a very virulent anaerobe that may cause severe infections, usually in children or young adults (1).

L. D. Medium or Lombard-Dowell Medium was developed by Dowell and Lombard(2) for the cultivation and identification of fastidious anaerobic bacteria. L. D. Esculin Agar is used to determine esculin hydrolysis, hydrogen sulfide and catalase production of *Bacteroides* and *Fusobacterium* species. L.D. Esculin HiVeg™ Agar is prepared by using vegetable peptones in place of animal based peptones which make the media free of BSE/TSE risks.

L. D. Agar is essentially a casein digest agar enriched with hemin, vitamin K1, L-cystine and yeast extract (3). This medium contains various nutritious substances, which can promote the growth of fastidious anaerobic bacteria. Tryptone and yeast extract provide the necessary nitrogenous nutrients while hemin and vitamin K1 supply additional growth factors. L-cystine and L-tryptophan serve as the amino acid sources. Esculin is hydrolyzed by the organisms to form esculetin and dextrose. The esculetin reacts with the iron salt of ferric citrate to produce a dark brown to black complex. Also L-cystine is a sulphur-containing amino acid and hence H<sub>2</sub>S production in combination with ferric citrate gives black colouration to the colonies. Vitamin K1 and hemin are the additional growth factors. Black colour of H<sub>2</sub>S positive colonies is rapidly lost after exposure to air; hence plates should be observed in anaerobic glove box or immediately upon exposure to air (3). Catalase-positive reaction may not be evident uptill 30 seconds to 1 minute after application of 3% hydrogen peroxide (3,4).

### Type of specimen

Water samples

### Specimen Collection and Handling:

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(5) After use, contaminated materials must be sterilized by autoclaving before discarding.

### Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection.

Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

### Limitations :

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

### Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 2.0% agar gel.

#### Colour and Clarity of prepared medium

Medium amber coloured clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 3.46% w/v aqueous solution at 25°C. pH : 7.4±0.2

#### pH

7.20-7.60

#### Cultural Response

Cultural characteristics observed under anaerobic condition after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	H <sub>2</sub> S production	Catalase	Esculin hydrolysis
<i>Bacteroides saccharolyticus</i> ATCC 25260	50-100	luxuriant	negative reaction	negative reaction	negative reaction
<i>Bacteroides fragilis</i> ATCC 25285	50-100	luxuriant	negative reaction	positive reaction, effervescence seen on addition of 3% hydrogen peroxide	positive reaction, brown black precipitate around the colonies
<i>Fusobacterium mortiferum</i> ATCC 9817	50-100	luxuriant	positive reaction, blackening of colonies	negative reaction	positive reaction, brown black precipitate around the colonies

### Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

### Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (6,7).

## Reference

1. Murray P. R., Baron J. H., Pfaller M. A., Tenover J. C. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
2. Dowell V. and Lombard G., June 1977, U.S., DHEW, Center for Disease Control (CDC), Atlanta. Ga.
3. Finegold S. M., Baron E. J., Bailey and Scotts Diagnostic Microbiology, 8th Ed., 1990, The C.V. Mosby Company.
4. Koneman E., Allen S., Dowell V. and Sommers H., 1979, Colour Atlas and Textbook of Diagnostic Microbiology, J. B. Lippincott Co., Philadelphia.
5. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
6. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
7. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 03/2024

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.