

Clostridium Difficile HiVeg™ Agar Base

MV836

Intended Use:

Recommended for selective isolation of *Clostridium difficile* from food and other specimens.

Composition**

Ingredients	g / L
HiVeg™ peptone No. 3	40.000
Disodium hydrogen phosphate	5.000
Potassium dihydrogen phosphate	1.000
Magnesium sulphate	0.100
Sodium chloride	2.000
Fructose	6.000
Agar	15.000
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 34.55 grams in 500 ml purified / distilled water. Heat gently to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add rehydrated contents of 1 vial of CC Selective Supplement I (FD010) together with 7% (v/v) defibrinated Horse blood or Sheep blood. Mix well and pour into sterile Petri plates.

Principle And Interpretation

The spectrum of disease caused by *Clostridium difficile* (a pathogenic *Clostridium* affecting the bowel) ranges from pseudomembranous colitis (PMC) through antibiotic associated colitis (AAC). It also includes chronic inflammatory bowel diseases, post-operative diarrhoea and non-antibiotic associated diarrhoea (1). Smith and King (2) first reported the presence of *C.difficile* in human infections. George et al (3) recommended the use of a fructose-containing medium with egg yolk for the isolation of *C.difficile* from faecal specimens. The medium was made inhibitory to the accompanying flora by the addition of the selective agents namely, D-cycloserine and cefoxitin.

HiVeg™ Agar Base w/ 1% Agar is prepared by completely replacing animal based peptone with vegetable peptones to avoid BSE/TSE risks associate with animal peptones. This medium does not contain neutral red indicator, as in the original formulation, as it is recommended for use with sheep or horse blood (3). Clostridium Difficile Agar Base is used for the primary isolation of *C.difficile* from faecal specimens. The medium composition is designed so as to obtain luxuriant growth of *C.difficile*. The selective agents D-cycloserine and cefoxitin used in the medium inhibit the growth of majority of *Enterobacteriaceae* and also *Enterococcus faecalis*, Staphylococci, gram-negative anaerobic bacilli and *Clostridium* species other than *C. difficile*, which may be found abundantly in faecal samples. Addition of 7% v/v horse blood to the base increases the recovery of *C. difficile* and also increases its colony size.

Spread a part of the faecal sample on the medium to obtain isolated colonies. Incubate the plates anaerobically at 37°C for 18 - 48 hours. *C. difficile* forms grayish white, irregular, raised and opaque colonies, 4-6 mm in diameter, after 48 hours. Typical gram stain morphology of *C. difficile* may not be seen in colonies taken from this medium due to the presence of antibiotics. Subculture on Blood Agar (M073) to obtain characteristic morphology. *C.difficile* colonies will not exhibit the typical fluorescence and colour of colony on this medium whereas other Clostridia can give fluorescence. Therefore, for complete identification and confirmation, other tests like gram staining, morphology, biochemicals, specific cytotoxin and clinical observation should be carried out.

Type of Specimen

Food samples.

Specimen Collection and Handling:

For food samples, follow appropriate techniques for sample collection and processing as per guidelines (4). After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Further biochemical and serological tests must be carried out for further identification.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Basal medium: Light amber coloured clear to slightly opalescent gel. After addition 7% v/v defibrinated horse blood: Cherry red coloured, opaque gel forms in Petri plates.

Reaction

Reaction of 6.91% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH

7.20-7.60

Cultural Response

Cultural characteristics observed under anaerobic condition with added CC Selective Supplement I (FD010) and 7% v/v defibrinated horse blood, after an incubation at 35-37°C for 48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Clostridium difficile</i> ATCC 11204	50-100	good-luxuriant	≥50%	greyish-white
<i>Shigella flexneri</i> ATCC 12022 (00126*)	≥10 ⁴	inhibited	0%	
<i>Escherichia coli</i> ATCC 25922 (00013*)	≥10 ⁴	inhibited	0%	
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923	≥10 ⁴	inhibited	0%	

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).

Reference

- Collee J. G., Fraser A. G., Marmion B. P., Simmons A., (Eds.), Mackie and McCartney, Practical Medical Microbiology, 14th Ed., Churchill Livingstone.
- Smith L. D. S. and King E. O., 1962, J. Bacteriol., 84:65.
- George W. L., Sutter V. L., Citron D., and Finegold S. M., 1979, J.Clin. Microbiol., 9:214
- Salfinger Y., and Tortorello M.L., 2015, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.

5.Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.

6.Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision :05/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.