

## Thiol HiVeg™ Broth

MV853

### Intended Use:

Recommended for cultivation of organisms from body fluids and other materials containing antibacterials such as Penicillin, Streptomycin and Sulphonamides.

### Composition\*\*

Ingredients	g / L
HiVeg™ peptone No.3	10.000
Yeast extract	5.000
Dextrose (Glucose)	1.000
Sodium chloride	5.000
Thiol compound	8.000
p-Amino benzoic acid (PABA)	0.050
Final pH ( at 25°C)	7.1±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 29.05 grams in 1000 ml purified/distilled water. Heat if necessary to dissolve the medium completely. Dispense in tubes or flasks to a depth of 6 cm for neutralization of Penicillin or in shallow layers for neutralization of Streptomycin. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

### Principle And Interpretation

Thiol Medium is used for culturing microorganisms from **body fluids** and also other materials containing antibiotics like penicillin, streptomycin or sulphonamides. The efficacy of Thiol Medium to retain viability of *Vibrio* was initially described by Huddleson (1). The ability of Thiol Medium to neutralize antibacterials was demonstrated by Christensen (2). This media can also be used for the cultivation and maintenance of *Haemophilus*, *Vibrio* and Meningococci (1). Thiol Broth which is Thiol Medium devoid of agar is also recommended for growing anaerobic bacteria in **blood** cultures and for recovery of nutritionally variant Streptococci (3,4) and *Bacteriodes* (5,6). Thiol HiVeg™ Broth is prepared by completely replacing animal based peptone with vegetable peptones to avoid BSE/TSE risks associate with animal peptones. HiVeg™ peptone No.3 and yeast extract provide nitrogenous compounds, vitamin B complex and other essential growth nutrients. Dextrose is the energy source. p-Amino benzoic acid serves as a preservative.

### Type of specimen

Please add specimens

### Specimen Collection and Handling:

After use, contaminated materials must be sterilized by autoclaving before discarding.

### Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

### Limitations :

1. Further biochemical and serological tests must be carried out for further identification.

### Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

### Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent solution.

### Reaction

Reaction of 2.9% w/v aqueous solution at 25°C. pH : 7.1±0.2

### pH

6.90-7.30

### Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18- 48 hours. Growth observed after addition of antibiotic concentrations upto 100 units of Penicillin or 1,000 micrograms of Streptomycin.

Organism	Inoculum (CFU)	Growth
<i>Neisseria meningitidis</i> ATCC 13090	50-100	poor-fair
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	good-luxuriant
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	good-luxuriant
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good-luxuriant

Key : \*Corresponding WDCM numbers.

### Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 15-25°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

### Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (7,8).

### Reference

1. Huddleson I. F., 1948, J. Bacteriol., 56:508.
2. Christensen D. H., 1947, Presented at the Michigan Branch, Society of American Bacteriologists, Detroit, Mich, December 12, 1947.
3. Donnelly J. P., 1994, Infect. Dis. Alert 6:109.
4. Isenberg (Ed.), 1992, Clinical Microbiology Procedures Handbook, Vol. 1, American Society for Microbiology, Washington, D.C.
5. Shanson D. C. and Barnicoat, 1975, J. Clin. Pathol., 28:407.
6. Szawatkowski M. V., 1976, Med. Lab. Sci., 33:5.
7. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
8. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 03/2024

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.