

B.T.B. Lactose HiVeg™ Agar

MV861

Intended Use:

Recommended for the detection and isolation of pathogenic Staphylococci.

Composition**

Ingredients	g / L
HiVeg™ peptone No. 3	5.000
HiVeg™ extract	3.000
Lactose	10.000
Bromo thymol blue	0.170
Agar	15.000
Final pH (at 25°C)	8.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 33.17 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Staphylococcus are known to be pathogenic to man and other mammals. Although this organism is frequently a part of the normal human microflora, it can cause significant opportunistic infections under appropriate conditions (1). Traditionally Staphylococci are divided into two groups on the basis of their ability to clot blood plasma (the coagulase reaction). The coagulase-positive Staphylococci constitute the most pathogenic species, *Staphylococcus aureus*. BTB Lactose Agar (2) designed by Chapman et al (3), is used in the detection and isolation of pathogenic Staphylococci. B.T.B. Lactose HiVeg™ Agar is prepared by completely replacing animal based peptone with vegetable peptones to avoid BSE/TSE risks associate with animal peptones. On this media Staphylococci are differentiated by their ability to grow at a high pH and in the presence of bromothymol blue. HiVeg™ peptone No. 3 and HiVeg™ extract serves as a source of carbon, nitrogen, long chain amino acids, vitamins and other essential nutrients. Lactose is the fermentable carbohydrate with bromothymol blue as an indicator. Plates should be inoculated preferably by spread plate technique and incubated for about 36 hours at 35°C. Typical colonies appear deep yellow (90% approx.) or blue grey (10% approx.). Coliforms may grow but are differentiated by their appearance.

Type of specimen

skin and mucous membranes

Specimen Collection and Handling

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

1. Individual organisms differ in their growth requirement and may show variable growth patterns on the medium.
2. Each lot of the medium has been tested for the organisms specified on the COA. It is recommended to users to validate the medium for any specific microorganism other than mentioned in the COA based on the user's unique requirement.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to greenish yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Greenish blue coloured, clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.32% w/v aqueous solution at 25°C. pH : 8.6±0.2

pH

8.40-8.80

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Escherichia coli</i> ATCC 25922 (00013*)	50-100	good-luxuriant	≥70%	yellow
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	good-luxuriant	≥70%	golden yellow
<i>Salmonella</i> Typhi ATCC 6539	50-100	good-luxuriant	≥70%	blue/colourless
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 6538 (00032*)	50-100	good-luxuriant	≥70%	golden yellow
<i>Staphylococcus epidermidis</i> ATCC 12228 (00036*)	50-100	good-luxuriant	≥70%	blue/colourless

Key : *Corresponding WDCM numbers.

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition. Seal the container tightly after use. Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (4,5).

Reference

1. Carney D. N., Fossieck B. E., Parker R. H. et al, 1982, Rev. Infect. Dis. H., 1-12.
2. Atlas R. M., 2004, Handbook of Microbiological Media, 3rd Edition, CRC Press.
3. Chapman, Lieb, Bereus and Curcio, 1937, J. Bacteriol., 33:533.
4. Isenberg, H.D. Clinical Microbiology Procedures Handbook 2nd Edition.
5. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.

Revision : 02/2024

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.