

Orchid Maintenance/ Replate Medium

With Calcium Chloride, Vitamins, Sucrose, Banana powder, HiVeg™ peptone, MES, Activated Charcoal and Agar

Product Code: PT074

Product Description :

Orchid Maintenance Medium has been formulated for the effective maintenance of the orchid species.

It is a nutrient blend of inorganic salts, vitamins and carbohydrates. In addition, it is supplemented with MES buffer which maintains optimum buffering and prevents acidification in the media required for the propagation of orchids. HiVeg™ peptone is added as an additional source of reduced organic nitrogen. Microelements like Boron, Manganese, Molybdenum, Copper, Iron and Zinc enhance metabolism in plants. Boron plays a key role in carbohydrate metabolism. Thiamine, pyridoxine, nicotinic acid act as enzymatic cofactors in universal pathways including glycolysis and TCA cycle and in primary and secondary metabolism in the plants. Activated charcoal adsorbs the inhibitory leachouts in the medium while banana powder aids in organogenesis as well as promotes the growth of protocorm like bodies.

The product is plant tissue culture tested but it is the sole responsibility of the user to ensure the suitability of the medium for individual species.

Composition :

Ingredients	mg/L
MACROELEMENTS	
Ammonium nitrate	825.000
Calcium chloride	166.100
Magnesium sulphate	90.340
Potassium nitrate	950.000
Potassium phosphate monobasic	85.000
MICROELEMENTS	
Boric acid	3.100
Cobalt chloride hexahydrate	0.013
Copper sulphate pentahydrate	0.013
EDTA disodium salt dihydrate	37.300
Ferrous sulphate heptahydrate	27.800
Manganese sulphate monohydrate	8.450
Molybdic acid (sodium salt)	0.106
Potassium Iodide	0.420
Zinc sulphate heptahydrate	5.300

VITAMINS

myo-Inositol	100.000
Nicotinic acid (free acid)	1.000
Pyridoxine HCl	1.000
Thiamine hydrochloride	10.000

CARBOHYDRATE

Sucrose	20000.000
---------	-----------

GELLING AGENT

Agar	7000.000
------	----------

OTHERS

Activated charcoal	2000.000
Banana Powder	30000.000
HiVeg™ Peptone	2000.000
MES	1000.000
Total(gms/litre)	64.3

Material required but not provided :

- Autoclaved distilled water
- Plant growth regulators
- 1N NaOH/HCl

Precautions :

- Ensure appropriate pH of the medium before addition of gelling agent as acidic pH will lead to decreased gelation resulting in semi solid flowing gel while alkaline pH will lead to formation of hardened gel.
- Use of Distilled water / Tissue culture grade water is recommended for media preparation as tap water or lower grade water may lead to salt precipitation and improper gelation.
- Avoid preparation of concentrated solutions, as it will lead to precipitation of salts.

Directions :

- Reconstitute medium by adding required quantity of powder in two-third of total volume with constant, gentle stirring till the medium gets completely dissolved.
- Add heat stable supplements prior to autoclaving.
- Make up the final volume with distilled water.
- Adjust the pH of the medium to 5.75 ± 0.5 using 1N NaOH/HCl.

- Heat the medium to boiling till complete dissolution of gelling agent.
- Sterilize the medium by autoclaving at 15 lbs and 121°C for 15 min.
- Cool the autoclaved medium to about 45°C before adding heat labile supplements.
- Aseptically dispense the desired amount of medium under a laminar airflow unit in sterile culture vessels.

Quality Control:

Appearance

Grey to black, homogenous, free flowing powder

Solubility

64.3 gms/litre soluble after boiling in distilled water

Colour and Clarity

Grey to black solution, opaque gel is formed on cooling

Gelling

Firm gel formed at pH : 5.75 ± 0.5

pH at 25°C

3.80 - 4.80

Plant Tissue Culture Test

The growth promoting properties of medium is assessed by providing plant cultures with relative humidity of about $60\% \pm 2\%$, temperature $22^\circ\text{C} \pm 2^\circ\text{C}$ and photoperiod of about 16:8. The plant species showed actively growing callus and shoots with no structural, necrotic and toxic deformity.

Storage and Shelf Life:

- The plant tissue culture medium powder is extremely hygroscopic and must be stored at 2-8°C in air tight containers.
- Preferably, entire content of each package should be used immediately after opening.
- Use before the expiry date.

Revision : 01 / 2017

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic , research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.