



Technical Datasheet

LoSera™ Nutrient Mixture F-10 Ham

With Sodium bicarbonate
Without L-Glutamine
1X Liquid Cell Culture Medium requiring reduced serum supplementation

Product Code: RSL009

Product Description:

LoSeraTM media are based on the classical formulations supplemented with insulin, transferrin and other advanced nutrients. The additional nutrients help in reducing the percentage of serum required to grow most of the common cell lines. The percentage of serum reduction may vary with type of cell line used. For nonfastidious cell lines serum can be reduced from 10% to as low as 1%. For fastidious cell lines serum usage can be reduced from 10% to 2.5%. LoSera™ medium can be used without prior adaptation and sub cultured using normal procedures. Reduced serum supplementation improves the reproducibility of experimental results by decreasing the variability caused due to undefined serum constituents. It also facilitates down regulation process in bioassays and in purification process of culture products.

RSL009 is LoSeraTM Nutrient Mixture F-10 Ham medium with Sodium bicarbonate. It does not contain L-glutamine. Users are advised to review the literature for recommendations regarding medium supplementation and physiological growth requirements specific for different cell lines.

Composition:

Ingredients	mg/L
INORGANIC SALTS	
Calcium chloride dihydrate	44.100
Copper sulphate pentahydrate	0.0025
Ferric sulphate heptahydrate	0.834
Magnesium sulphate anhydrous	74.730
Potassium chloride	285.000
Potassium dihydrogen phosphate	83.000
Sodium bicarbonate	1200.000
Sodium chloride	7400.000
Sodium phosphate dibasic anhydrous	153.700
Zinc sulphate heptahydrate	0.0288
AMINO ACIDS	
Glycine	7.510
L-Alanine	8.910
L-Arginine hydrochloride	211.000

L-Asparagine monohydate	15.010
L-Aspartic acid	13.300
L-Cysteine hydrochloride monohydrate	35.130
L-Glutamic acid	14.700
L-Histidine hydrochloride monohydrate	21.000
L-Isoleucine	2.600
L-Leucine	13.100
L-Lysine hydrochloride	29.300
L-Methionine	4.480
L-Phenylalanine	4.960
L-Proline	11.500
L-Serine	10.500
L-Threonine	3.570
L-Tryptophan	0.600
L-Tyrosine disodium salt dihydrate	2.610
L-Valine	3.500
VITAMINS	0.024
Biotin	0.698
Choline chloride	0.715
D-Ca-Pantothenate	1.320
Folic acid	0.615
Nicotinamide	0.206
Pyridoxine hydrochloride	0.376
Riboflavin	1.000
Thiamine hydrochloride	1.360
Vitamin B12	0.541
i-Inositol	0.5 11
OTHERS D-Glucose	1100.000
	Proprietary
Growth Supplement mix Hypoxanthine sodium salt	4.080
* ±	0.210
Lipoic acid Phenol red sodium salt	1.300
	110.000
Sodium pyruvate	0.730
Thymidine	0.750

Directions:

1. Add 20ml of 200mM L-glutamine (TCL012) or HiGlutaXL $^{\text{TM}}$ supplement (TCL030) for 1 litre of medium.

Recommendations for use with LoseraTM Media:

- 1. LoSeraTM media have been optimized at 2.5% serum concentration for a broad range of cell culture applications. Recommended concentrations of serum using LoSeraTM media ranges from 1-5%. However the concentration of serum used may need to be adjusted for specific cell types or applications to achieve optimal results. Titration of FBS concentration is recommended to determine maximum serum reduction.
- 2. LoSera[™] media are provided as 1X solutions and need to be supplemented with 4mM Glutamine and required amount of reduced serum.
- 3. In case of antibiotics being used to control contamination, it is recommended to reduce the amount of antibiotics in proportion to the amount of serum reduced.

Material required but not provided:

L-Glutamine Solution 200mM (TCL012) HiGlutaXL™ Supplement (TCL030) Fetal Bovine Serum (RM1112/RM10432)

Quality Control:

Appearance

Orangish red colored, clear solution.

pН

7.00 - 7.60

Osmolality in mOsm/Kg H2O

285.00 -325.00

Sterility

No bacterial or fungal growth is observed after 14 days of incubation, as per USP specification.

Cultural Response

The growth promotion capacity of the medium is assessed qualitatively by analyzing the cells for the morphology and quantitatively by estimating the cell counts and comparing it with a control medium.

Storage and Shelf Life:

Store at 2-8°C away from bright light. Shelf life is 12 months. Use before expiry date given on the product label.

Disclaimer: Revision: 03/2022

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

