



## Nystatin NS 100 Units

# **SD025**

Nystatin NS 100 Units discs are used for antimicrobial susceptibility testing of fungal cultures

## Composition

*Ingredients	Concentration
Nystatin	100 Units/disc

# **Susceptibility Test Procedure:**

Preparation of Inoculum:

- 1. Inoculum is prepared by picking five distinct colonies of approximately 1mm from 24 hours old culture grown on Sabouraud Dextrose Agar (M063) and incubated at  $35 \pm 2^{\circ}$ C. Colonies are suspended in 5ml of sterile 0.85% Saline.
- 2. Vortex the resulting suspension and adjust the turbidity to yield  $1 \times 10^6$  5 x  $10^6$  cells /ml (i.e. 0.5 McFarland standard).

Test Procedure:

- 1. Prepare plates with Muller Hinton Agar + 2% Glucose + 0.5 mcg/ml Methylene Blue Dye (M1825) for carrying out susceptibility of antifungal discs. The medium in the plates should be sterile and have a depth of about 4 mm.
- 2. Dip a sterile non-toxic cotton swab on a wooden applicator into the standardized inoculum (turbidity so adjusted, as to obtain semi confluent growth on the petri plate) and rotate the soaked swab firmly against the upper inside wall of the tube to express excess fluid. Streak the entire agar surface of the plate with the swab three times, turning the plate at 60° angle between each streaking. Allow the inoculum to dry for 5 15 minutes with lid in place.
- 3. Apply the discs using aseptic technique. Deposit the discs with centers at least 24 mm apart.
- 4. Invert the plates and place in an incubator set to  $35 \pm 2^{\circ}$ C within 15 minutes after the discs are applied.
- 5. Examine each plate after 20 24 hours of incubation. If plate was satisfactorily streaked the resulting zones of inhibition will be uniformly circular and there will be a semi-confluent lawn of growth. Read at 48 hours only when insufficient growth is observed after 24 hours incubation.

### **Principle:**

Antimicrobial susceptibility testing (AST) of bacterial and fungal isolates is a common and important technique in most clinical laboratories. The results of these tests are used for selection of the most appropriate antimicrobial agent(s) for treatment against the infectious organisms. Till the 1950s, laboratories were lacking in the methodologies and equipments for the accurate determination of in vitro responses of organisms to antimicrobial agents. Bauer et al (1) began the development of standardized methods for antimicrobial susceptibility testing, using disc diffusion system. However the susceptibility results may not always correlate with the patient's response to therapy. The response of an infected patient to antimicrobial agent(s) is a complex interrelationship of host responses, drug dynamics and microbial activity. Antimicrobial susceptibility tests are either quantitative or qualitative. Disc diffusion test is a qualitative test method. The National Committee for Clinical Laboratory Standards (NCCLS), now known as Clinical Laboratory Standards Institute (CLSI) has published comprehensive documents regarding the disc diffusion systems. The agar disc diffusion test is the most convenient and widely used method for routine antimicrobial susceptibility testing. In subsequent and current practice, antimicrobial impregnated paper discs are applied onto the agar surface. Based on the Bauer-Kirby Method, standardized reference procedures for the disc systems were published by WHO and FDA and are periodically updated by the CLSI (formerly NCCLS)(2). For any antimicrobial testing, Quality control or clinical testing, the method to be followed is the same as mentioned above.

However few precautions are to be maintained while handling of the Sensitivity discs,

- On receipt the discs are to be immediately stored at the recommended temperature.
- Medium preparation, Inoculum preparation and incubation to be done as specified.

## **Quality Control:**

Appearance: Filter paper discs of 6mm diameter with printed "NS 100" on centre of each side of the disc.

**Cultural response**: Average diameter of zone of inhibition observed on Muller Hinton Agar + 2% Glucose + 0.5 mcg/ml Methylene Blue Dye after 24-48 hours incubation at 35-37°C for standard cultures.

Organisms (ATCC)	Std. zone of diameter (mm)
<i>C.albicans</i> (90028)*	19-27
C.parapsilosis (22019)*	16-25
C.tropicalis (750)*	16-21
C.krusei(6528)*	15-20
C.albicans(10231)	15-23
S.cerevisiae(9763)	17-25

\* = Q.C. Strains recommended by CLSI

#### **Storage and Shelf-life:**

Discs should always be stored at  $-20^{\circ}$ C to  $+8^{\circ}$ C under dry conditions, along with the dessicator provided in individual pack. Use before expiry date on the label.

#### **References:**

1. Method for Antifungal Disk Diffusion Susceptibility Testing of Yeasts; Approved Guidelines-Second edition Vol.29 No.17, August- 2009 CLSI document M44-A2. For more details refer to this volume

Note :

Use following media to carry out susceptibility test For rapidly growing aerobic organisms : Mueller Hinton Agar (M173/M1084) For *Haemophilus* spps : Haemophilus Test Agar (M1259 + FD117) For *S.pneumoniae* : Muller Hinton Agar supplemented with 5% Sheep Blood For Neisseria spps : G.C.Agar +1% defined growth supplement (M434 + FD025)

#### \* Not for Medicinal Use

IVD	In vitro diagnostic medical device
Œ	CE Marking



Do not use if package is damaged

Storage temperature

HiMedia Laboratories Pvt. Limited, C-40, Road No.21Y, MIDC, Wagle Industrial Area, Thane (W) -400604, MS, India

REP EC

CE Partner 4U, Esdoornlaan 13, 3951 DB Maarn The Netherlands, <u>www.cepartner</u> 4u.eu

#### Disclaimer :

Revision : 03 / 2022

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia<sup>™</sup> publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia<sup>™</sup> Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.