

L-Histidine monohydrochloride, monohydrate

(From non-animal source)
Cell Culture Tested

Product Code: TC077

Product Description :

Molecular Weight: 209. 6

Molecular Formula: $C_6H_9N_3O_2 \cdot HCl \cdot H_2O$

CAS No.: 5934-29-2

Synonym: (S)-a-amino-1H-imidazole-4-propanoic acid hydrochloride, glyoxaline-5 alanine hydrochloride, His, H

L-Histidine monohydrochloride monohydrate is chloride salt of L-Histidine. L-Histidine is positively charged hydrophilic, essential α -amino acid coded by codons CAU and CAC. It is chemically basic in nature. It carries positively charged imidazole functional group.

It is used as a major component in wide range of cell culture media including classical and serum-free media. It plays many important roles in cell culture as L-Histidine. Some of them are mentioned below:

1. Protein synthesis and protein folding:

Like all other amino acids, L-Histidine also acts as a substrate for protein synthesis during translation process. Histidine's side chain allows it to act as both a base and an acid, both donating and accepting protons, which can be of considerable importance in its role as part of proteins. Hence it is a proteinogenic amino acid.

2. Nucleic acid synthesis:

Biosynthesis of Histidine is inherently linked to the pathways of nucleotide formation. Thus it participates in synthesis of nucleic acids also.

3. As a catalyzer:

Because of presence of imidazole ring, Histidine is a nucleophile and a good acid/base catalyzer. Histidine residues are found in enzyme active sites.

4. Precursor for synthesis of carnosine:

Histidine acts as a precursor for synthesis of carnosine, a dipeptide of amino acids beta-alanine and histidine. Carnosine exerts anti-oxidant effect and protects cellular proteins by preventing oxidation of sugars. It also binds with potentially harmful carbonyl groups that attack and bind with proteins imbedded in cell membrane.

Directions :

Preparation instructions:

L-Histidine hydrochloride monohydrate is soluble in water (100mg/ml).

Quality Control:

Appearance

White to offwhite crystalline powder.

Solubility

Clear colorless solution at 10gm in 100ml of water .

pH of 10% solution in water

3.50 -4.50

Specific rotation $[\alpha]_{20/D}$

+8.9° to +9.5°

Chloride (Cl)

NMT 0.02%

Ammonium (NH₄)

NMT 0.02%

Sulphate (SO₄)

NMT 0.02%

Iron (Fe)

NMT 0.001%

Heavy metals

NMT 0.001%

Arsenic (As)

NMT 0.0001%

Loss on drying

NMT 0.2%

Assay

NLT 99%

Cell Culture Test

Passes

Storage and Shelf Life:

Store at 10-30°C away from bright light.

Shelf life is 48 months.

Use before expiry date given on the product label.

Revision : 1 / 2011

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.