

# HiKaryoXL™ Colchicine Solution

With 10µg per ml Colchicine in phosphate buffered saline

**Product Code: TCL062**

## Product Description:


Colchicine is an alkaloid that is isolated from the plant *Colchicum autumnale*. Colchicine is used for arresting the cells in metaphase. Arresting of cells in metaphase allows an increased yield of mitotic cells for analysis. Colchicine inhibits microtubule polymerisation by binding to tubulin. Availability of tubulin is essential for mitosis and therefore colchicine effectively arrest cells in metaphase.

TCL062 is a sterile filtered solution of 10µg per ml Colchicine in phosphate buffered saline.

## Directions:

### For Peripheral Blood Culture

1. Add 500µl freshly collected heparinized whole blood to 5ml of HiKaryoXL™ Medium in a sterile 15ml conical bottom centrifuge tube.
2. Loosen the cap of tube by one thread and incubate at 37°C and 5% CO<sub>2</sub> for 70 - 72 hours in horizontal position as shown in the figure below:



*Note: Alternatively, the tubes can be incubated in a non-CO<sub>2</sub> incubator. Absence of CO<sub>2</sub> does not affect the mitotic count.*


3. Add 250µl of 10µg/ml of Colchicine (TCL062) and incubate at 37°C for additional 20 minutes.

*Note: Incubation of 2 hours gives higher mitotic count than 20 minutes. Users are advised to decide incubation time as per their need and convenience.*

4. After incubation with colchicine, centrifuge the tubes at 2000rpm for 4 minutes.
5. Discard the supernatant and vortex briefly for 5 seconds to disperse the pellet uniformly.
6. Add 5ml 0.075M Potassium chloride solution (TCL040) and incubate at room temperature for 7 minutes keeping the tubes in an upright position. Mix by inverting.
7. Centrifuge the tubes at 2000rpm for 4 minutes.
8. Discard the supernatant and vortex briefly for 5 seconds to disperse the pellet uniformly.
9. Add 5ml of freshly prepared ice cold fixative drop by drop (Acetic acid: methanol, 1:3 parts) and mix gently by inverting.

*Note: Addition of fixative for the first time may create turbulence which in turn may lead to cell breakage and irreversible clumping. Hence, fixative addition for the first time should be done dropwise and slowly.*

10. Repeat steps 7, 8 and 9 two more times.
11. Resuspend the pellet in 0.5ml of fresh fixative and store them at -20°C till slide preparation.
12. Clean the slides with mild detergent and wash thoroughly under tap water to make them grease free.
13. Place the clean slides in a beaker containing water such that they are completely immersed in water. Keep the beaker in a refrigerator at 2 – 8°C and allow the slides to cool. *Note: Steps 12 and 13 can be performed during incubation period of 2-4 hours with colchicine solution to save time.*
14. Mix the cell suspension gently by pipetting up and down. DO NOT vortex.
15. Hold the ice cold wet slide at 45° angle and drop 50µl suspension at the bottom of slide with the help of micropipette in such a way that the suspension hits hard on the slide and then runs down surface. Refer the figure mentioned below:



### Storage and Shelf Life:

Store at 2-8°C.

Shelf life of the product is 12 months.

Use before expiry date given on the product label.

16. Similarly drop 50µl suspension the center and 50µl at the top of the slide.

*Note: Ensure that the direction of dropping is from bottom to the top.*

17. Allow the slides to air dry. DO NOT blow.
18. Heat fix the slides by holding them over a hot plate for 10 – 12 seconds, with chromosome spreads facing up.
19. Stain the slides with required staining solution.

### Material required but not provided:

HiKaryoXL™ RPMI Medium (AL165A) or

HiKaryoXL™ Nutrient Mixture F-10 Ham Medium (AL169A)

Potassium Chloride solution 0.075M (TCL040)

Methanol

Acetic Acid

Giemsa Stain (TCL083)

### Quality Control:

#### Appearance

Colorless, clear solution.

#### Sterility

No bacterial or fungal growth is observed after 14 days of incubation, as per USP specification.

#### Performance Tests

##### 1. Peripheral blood performance test

Passes

##### 2. Mitotic index (Test lot)

NLT 0.010

##### 3. Mitotic index (Control lot)

NLT 0.010

### Disclaimer :

Revision : 04/2025

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic , research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.