

Woody Plant Macroelements

TS1036

Composition :

Ingredients	milligrams/litre
Ammonium nitrate	400.00
Calcium chloride.2H ₂ O	96.00
Calcium nitrate anhydrous	386.34
Magnesium sulphate	180.69
Potassium phosphate monobasic	170.00
Potassium sulphate	990.00
TOTAL gm/litre	2.22

Directions :

Suspend 2.20 grams of dehydrated macroelements powder[#] in 600ml of distilled water. Apply constant gentle stirring to the solution till the powder dissolves completely. Add desired heat stable supplements prior to autoclaving. Adjust the medium to the desired pH using 1N HCl/NaOH. Make up the final volume to 1000ml with distilled water. Sterilize the medium by autoclaving at 15 lbs or 121°C for 15 minutes. Cool the autoclaved medium to 45°C before adding the filter sterilized heat labile supplements. Dispense the desired amount of medium aseptically in sterile culture vessels.

Weight after vacuum drying to remove all water

Principle and Interpretation :

Woody plant macroelements has been specially formulated for plant cell, tissue and organ cultures. Ammonium nitrate and calcium nitrate serves as the sources of nitrate.

Quality Control :

Appearance	: White to off-white, homogeneous, free flowing powder.
Solubility	: 2.20 gm/litre soluble in distilled water.
Colour and Clarity	: Colourless to light yellow, clear solution.
pH at 25°C	: 4.9 \pm 0.5 of 0.220% w/v dehydrated macroelements powder.

Cultural Response :

Cultural condition :

 Incubation p Relative hur Temperature Photoperiod 	nidity	: 5 weeks : 60% ± 2% : 22°C ± 2°C : 16:8
Cell Line	Type of Culture	Results
Musa species	Shoot culture	No structural deformity observed
		No necrotic tissues,
		Actively growing shoots,
		No toxicity to shoots
Daucus species	Callus culture	No necrotic tissues,
		Actively growing callus,
		No toxicity to callus

[The medium is prepared as per direction. The growth promoting activity of this dehydrated macroelements is evaluated using two plant species viz. Musa species and Daucus species through three passages. Plant growth hormones (e.g. 2,4-D, NAA, Kinetin and 6-BAP) are added in suitable combinations and concentrations.]

Storage and shelf life :

Dehydrated macroelements powder is extremely hygroscopic and should be protected from atmospheric moisture. If possible, the entire content of each bottle should be used immediately after opening or else the unused portion should be stored in a desiccator and refrigerated at 2-8°C. Use before the expiry date.

Reference :

1. Lloyd G. & McCown B., Int. Plant Prop. Soc. Proc., (1981), 30, 421 - 427

Disclaimer :

CE

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia[™] publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia[™] Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

HiMedia Laboratories Pvt. Ltd. A-516, Swastik Disha Business Park, Via Vadhani Ind. Est., LBS Marg, Mumbai-400086, India. Customer care No.: 022-6147 1919 Email: techhelp@himedialabs.com